

Hidden Quantum Gravity in 3d Feynman diagrams

Aristide Baratin¹, Laurent Freidel^{1,2}

¹*Laboratoire de Physique, École Normale Supérieure de Lyon*

46 allée d'Italie, 69364 Lyon Cedex 07, France.

²*Perimeter Institute for Theoretical Physics*

31 Caroline street North, N2L2Y5 Waterloo, ON, Canada

Abstract

In this work we show that 3d Feynman amplitudes of standard QFT in flat and homogeneous space can be naturally expressed as expectation values of a specific topological spin foam model. The main interest of the paper is to set up a framework which gives a background independent perspective on usual field theories and can also be applied in higher dimensions. We also show that this Feynman graph spin foam model, which encodes the geometry of flat space-time, can be purely expressed in terms of algebraic data associated with the Poincaré group. This spin foam model turns out to be the spin foam quantization of a BF theory based on the Poincaré group, and as such is related to a quantization of 3d gravity in the limit $G_N \rightarrow 0$. We investigate the 4d case in a companion paper [1] where the strategy proposed here leads to similar results.

Contents	
I. Introduction	3
II. Feynman diagrams in terms of invariant measure	4
III. Feynman diagrams as spin foam models	7
A. The main result	7
B. Emergence of the model	8
1. A key identity	8
2. Spin foam in Feynman graph: inducing the geometry dynamically	11
C. Topological invariance	11
D. Symmetries and gauge fixing	14
E. Observables and partial gauge fixing	17
IV. Algebraic structure	19
A. A Poincaré model	19
1. The result	19
2. Proof of the Theorem	20
B. Poincaré versus Ponzano-Regge	24
V. Feynman diagrams on homogeneous spaces	25
VI. Gravity and BF theory	26
VII. Conclusion	27
A. Geometry of simplicial complexes	28
1. Basic definitions and discussion about orientation	28
2. On the geometry of the simplex	31
B. Flatness identities for the tetrahedron	33
References	35

I. INTRODUCTION

There has been a recent understanding and new results concerning the consistent coupling of matter fields to 3d quantum gravity amplitudes, in the context of the spin foam approach to 3d quantum gravity [2, 3, 4, 5, 6, 7]. These results have led to the construction of an effective field theory, describing the coupling of quantum fields to quantum gravity [8, 9], which arises after the exact integration of quantum gravitational degrees of freedom. This effective field theory lives on a non-commutative space-time whose geometry is studied in detail in [10]. When the no gravity limit $G_N \rightarrow 0$ is taken, the geometry becomes commutative and usual field theory is recovered.

From the spin foam side, the way usual Feynman diagrams are recovered is highly non trivial, since quantum gravity amplitudes are expressed as purely algebraic objects in terms of spin foam models, a language quite remote from usual field theory language. The fact that Feynman diagrams arise from this picture effectively, amounts to show that usual field theory can be expressed in a background independent manner, where flat space-time emerges dynamically from the choice of spin foam amplitudes. Namely, this amounts to show that Feynman amplitudes can be expressed as the expectation value of certain observables in a topological spin foam model.

Note that this last statement is not a statement regarding quantum gravity but one regarding well known Feynman diagrams in 3d. It is then natural to wonder if one can reach a deeper understanding of this property; whether it is tied up to the topological nature of gravity in dimension 3 or if it can be extended to field theory in higher dimensions. It also begs the question, whether or not Feynman diagrams contain some seeds of information about quantum gravity dynamics.

A natural strategy to address successfully these questions is to start from a careful study of the structure of Feynman graph amplitudes, without any assumption about the nature of quantum gravity dynamics. The goal is to see why and how such familiar objects can be interpreted in terms of spin foam models, which have appeared to be a suitable language to address dynamical question in a background independent approach to quantum gravity [11].

Usual Feynman diagrams are natural observables¹ from which physical predictions regarding matter interactions in space-time can be extracted. As such, they allow one in principle to probe the geometry of space-time at arbitrarily small scales. Moreover, they can be naturally promoted to physical observables in any theory of quantum gravity. Therefore, the preferred spin foam model - if any - which is related to Feynman diagrams should give us some important seeds of information about the structure of quantum gravity amplitudes. In other words, if one assumes that it is possible to consistently describe quantum gravity amplitudes coupled to matter in terms of a spin foam model, a necessary requirement for the spin foam amplitudes, in order to be physically relevant, is to reduce to the ones appearing in the study of Feynman diagrams when $G_N \rightarrow 0$. This gives strong restrictions on the admissible, physically viable spin foam models.

In this paper we focus our study to the case of 3-dimensional Feynman diagrams, but as should be clear from our work and as shown in a companion paper [1], most of the results obtained here can be extended to the more interesting and challenging case of 4-dimensional amplitudes. What makes the analysis work is not that gravity is a topological theory but the fact that, when coupled to matter, quantum gravity *in the limit* $G_N \rightarrow 0$ behaves as a topological field theory. This is a trivial fact in dimension 3 and has been argued to be true in dimension 4 when one takes this limit while preserving diffeomorphism invariance [12].

The main idea behind our analysis is the fact that in 3d, quantum gravity amplitudes provide an

¹ Of course, any observable quantity is actually given by a sum of Feynman diagrams, for instance those of fixed valency and given degree, as generated by a field theory at given order of perturbation, and subject to renormalization procedure. In order to keep the exposition simple we will refer to individual Feynman diagrams, in a regularized form in order to avoid divergences.

integration measure for the integration of Feynman graph amplitudes. This idea is contained in [8] and spelled out in [13]. When gravity is turned off, this measure should be related to the Lebesgue measure expressed in terms of relative lengths, which are Poincaré scalars. When gravity is turned on, this measure is deformed and reveals the quantum nature of the corresponding space-time.

The organization of this paper is as follows. In section II we first express explicitly the change of variables from the Lebesgue measure that arises in Feynman graph integration, to the measure expressed in terms of relative distances. In section III we present our main result, namely that Feynman integrals can be expressed as the expectation value of an observable for a topological spin foam model. This model is constructed out by removing all explicit information about flat space into a uniquely defined choice of amplitudes. We carefully study this model, especially its symmetries and its gauge fixing properties, and show that it defines a topological state sum model. This topological model, which naturally emerges from Feynman graphs, has already been studied in the mathematical literature from a different perspective by Korepanov [14]. In section IV we show that the tetrahedral weights entering the definition of the topological state sum model can be given an algebraic interpretation in terms of the $6j$ -symbol of the Poincaré group; this $6j$ -symbol is computed explicitly and for the first time. We also relate this model to the Ponzano-Regge model via a doubling and limiting procedure. Finally we show how to extend these results to the case of a non vanishing cosmological constant.

II. FEYNMAN DIAGRAMS IN TERMS OF INVARIANT MEASURE

We consider QFT in flat Euclidean D -dimensional space-time and focus on closed Feynman diagrams for a scalar field. The Feynman amplitude for a graph Γ is given by an integral over the position in $\mathbb{R}^D$ of N vertices. The (Euclidean) Poincaré group ISO_D acts diagonally on these N points

$$(\vec{x}_1, \dots, \vec{x}_N) \rightarrow (\Lambda \cdot \vec{x}_1 + \vec{a}, \dots, \Lambda \cdot \vec{x}_N + \vec{a})$$

where $(\Lambda, \vec{a}) \in SO_D \times \mathbb{R}^D$ is a general element of the Euclidean Poincaré group. The integrand is a function of the distances between the vertices, namely an invariant under the action of ISO_D

$$I_\Gamma = \int_{\mathbb{R}^D} d^D x_1 \cdots d^D x_N \mathcal{O}_\Gamma(|\vec{x}_i - \vec{x}_j|), \quad \text{with} \quad \mathcal{O}_\Gamma = \prod_{(ij) \in \Gamma} G^F(\vec{x}_i - \vec{x}_j) \quad (1)$$

G_F is the Feynman propagator and the product is over all edges of Γ . Following the ideas of [8, 13], we want to express this integral in terms of the invariant measure, involving the quantities $l_{ij} = |\vec{x}_i - \vec{x}_j|$. On one hand, ND variables are integrated over in (1) while the dimension of the Poincaré group is $\frac{1}{2}D(D+1)$; consequently, the invariant measure depends on $ND - \frac{1}{2}D(D+1)$ variables: the number of independent variables required to place the vertices with respect to each other². On the other hand, the number of length variables l_{ij} is $\frac{1}{2}N(N-1)$. Note that the quantity:

$$\frac{N(N-1)}{2} - \left[ND - \frac{D(D+1)}{2} \right] = \frac{(N-D)(N-D-1)}{2} \quad (2)$$

which computes the number of redundant edge lengths, vanishes only for $N = D$ and $N = D + 1$.

² Precisely this quantity is the number of *continuous* variables required to place the vertices with respect to each other modulo *discrete* transformations.

First, for $N = D$, we split the measure into the Haar measure for ISO_D and a product of $l_{ij}dl_{ij}$

$$d^D x_1 \cdots d^D x_D = d^D a d\Lambda \prod_{i < j} l_{ij} dl_{ij}, \quad (3)$$

where a is the barycenter of the D points and $d\Lambda$ is the Haar measure on SO_D . The volume of SO_D with respect to this measure is given by

$$A_D = \frac{2^{D-1} \pi^{\frac{D(D+1)}{4}}}{\prod_{k=1}^D \Gamma(\frac{k}{2})},$$

so $A_2 = 2\pi$, $A_3 = 2^3\pi^2$ and $A_4 = 2^4\pi^4$. Given an additional point x_{D+1} , one can build a D -simplex in $\mathbb{R}^D$ with vertices x_i , and express the Lebesgue measure in terms of the edge lengths as

$$d^D x_{D+1} = \sum_{\epsilon = \pm 1} \frac{\prod_{i=1}^D l_{iD+1} dl_{iD+1}}{\mathcal{V}(l_{ij})} \quad (4)$$

$\mathcal{V}(l_{ij})$ is $D!$ times the volume of the simplex whose edge lengths are l_{ij} . In the rest of the paper, we will simply call ‘volume’ the quantity $\mathcal{V}$. ϵ labels the orientation³ of the D -simplex $x_1 \cdots x_{D+1}$.

The proof of (4) is easily obtained by computing the jacobian of the linear function which maps an orthonormal basis e_i to the basis defined by the vectors $\{x_i\}_{i=1 \dots D}$. Let $y \in \mathbb{R}^D$ and $l_i = |y - x_i|$; we have

$$\prod_{i=1}^D l_i dl_i = \prod_{i=1}^D d \frac{(y - x_i)^2}{2} = \prod_{i=1}^D dy_j \det \left(\frac{\partial}{\partial y^j} \frac{(y - x_i)^2}{2} \right) \quad (5)$$

$$= \prod_{i=1}^D dy_i |\det(e_j \cdot (y - x_i))| = d^D y \sqrt{\det((y - x_j) \cdot (y - x_i))} \quad (6)$$

This last determinant is $D!$ times the volume of the simplex $x_1, \dots, x_D, y$. The previous identity is a local identity showing that the map $y \rightarrow l_i$ is locally invertible away from the hyperplane spanned by $x_1, \dots, x_D$. This map is globally invertible on each half space separated by this hyperplane. The sum over orientations comes from the fact that the set of edge lengths l_i determines the position of y up to a reflection with respect to the hyperplane spanned by $x_1, \dots, x_D$. Such a reflection is in $O(D)$, but not in $SO(D)$, and changes the orientation of the D -simplex. The equality of measures (4) is then understood as an equality for integrating out a function of x_{D+1} in the LHS and a function of l_{iD+1} and orientation ϵ in the RHS. The proof of (3) is obtained by a direct computation and by recurrence.

So putting (3,4) together we get, for $N = D + 1$ points in $\mathbb{R}^D$, that the invariant measure is given by:

$$d^D x_1 \cdots d^D x_{D+1} = d^D a d\Lambda \sum_{\epsilon} \frac{\prod_{i,j} l_{ij} dl_{ij}}{\mathcal{V}(l_{ij})}. \quad (7)$$

A similar identity already appeared in [15].

Now for $N = D + 2$, the quantity (2) is equal to 1. Accordingly, the invariant measure will be formulated using all the distances, except for one. One can indeed express it in terms of the edge lengths of the polyhedron formed by two D -simplices glued together in $\mathbb{R}^D$ along a $(D - 1)$ -simplex

³ The notion of orientation is precisely defined in appendix A.

(as shown in Fig 2 for $D = 3$). The missing edge length is the one joining the only two vertices, denoted by a, b , which do not belong to the common $(D - 1)$ -face. The measure is obtained by using twice (4)

$$d^D x_1 \cdots d^D x_{D+2} = d^D a d\Lambda \sum_{\epsilon_a, \epsilon_b} \frac{\prod_{(ij) \neq (ab)} l_{ij} dl_{ij}}{\mathcal{V}_a \mathcal{V}_b} \quad (8)$$

where the product is over all edges (ij) different from (a, b) , $\mathcal{V}_a$ denotes the volume of the D -simplex not containing a , and the sum is over orientations of the D -simplices a, b .

Note that the edge lengths $l_{ij}, (ij) \neq (ab)$ only determine the geometry of the polyhedron modulo reflections of the D -simplices along the common $(D - 1)$ -face. In order to fully determine the geometry, one needs to specify the orientations of the two D -simplices denoted a and b . Once these orientations ϵ_a, ϵ_b are specified, the missing edge length can be uniquely reconstructed: $l_{ab} = l_{ab}(l_{ij}, \epsilon)$. For fixed l_{ij} , the missing distance l_{ab} can take two values denoted by $l_{ab}^\pm$, with $l_{ab}^- \leq l_{ab}^+$, depending on whether the two points (a, b) are separated or not by the hyperplane spanned by the common face. A simultaneous change of the two orientations is equivalent to a symmetry with respect to this hyperplane, under which all distances remain invariant. Therefore, the length l_{ab} depends only on the *product* $\epsilon_a \epsilon_b$, in such a way, according to our conventions, that $l_{ab}^\pm = l_{ab}^{\epsilon_a \epsilon_b}$. The Fig 1 illustrates the $D = 2$ case. More details are given in appendix.




FIG. 1: Two triangles acd and bcd in the Euclidean plane share an edge (cd) . On the left (resp. right) $l_{ab} = l_{ab}^+$ (resp. l_{ab}^-).

For a general value of $N = D + k$, we use repeatedly (k times) the equation (4) in order to get the invariant measure. This invariant measure is based on a triangulation which is constructed recursively: each time a vertex is added, we choose a $(D - 1)$ -face on the existing triangulation, and build a new D -simplex from this data. At each step we use (4) to construct the invariant measure whose form depends on the triangulation Δ_k . The triangulations obtained this way are of special type: they are triangulations with a boundary being topologically a $(D - 1)$ -sphere and such that every $(D - 2)$ -face lies on the boundary.

More formally, let B_D be a D -dimensional ball B_D with $n = D + k$ points marked on its boundary. We now consider a triangulation Δ_k of B_D with $D + k$ vertices, such that every vertex and every $(D - 2)$ -face are on the boundary. In other words Δ_k is a triangulation of B_D which possesses $D + k$ boundary vertices and which does not possess any internal $(D - 2)$ -face. If one draws the dual 2-skeleton of Δ_k , in which vertices are dual to D -simplices, edges to $(D - 1)$ -simplices and faces to $(D - 2)$ -simplices, one sees that our condition on Δ_k implies that this dual graph is a D -tree (a $D + 1$ valent graph containing no closed loop) with open ends. It can be checked that the number of edges of such a triangulation matches the number of Poincaré invariant quantities.

Given such a triangulation the invariant measure reads

$$d^D x_1 \cdots d^D x_{D+k} = d^D \text{ad} \Lambda \sum_{\epsilon \in \{\pm 1\}^k} \prod_{e \in \Delta_k} l_e d l_e \prod_{\sigma \in \Delta_k} \frac{1}{\mathcal{V}_\sigma} \quad (9)$$

where the first product is over all edges e of the triangulation Δ_k , the second product is over all its D -simplices σ and the summation is over all orientations of its D -simplices.

By construction the data $\{l_e, \epsilon_\sigma\}$ defines uniquely a flat geometry on Δ_k and allow us to reconstruct the ‘missing’ edge lengths, namely the length of the edges that do not belong to Δ_k . The set of distances $\{l_{ij}\}$ in the integrand of (1) splits up into the lengths $\{l_e, e \in \Delta_k\}$ and the ‘missing’ distances denoted by $\{l_{e'}^\epsilon\}$. These missing distances are ϵ -dependent functions of the l_e .

The Feynman amplitude (1) is finally given by:

$$I_\Gamma = \int \prod_{e \in \Delta_k} l_e d l_e \sum_{\epsilon \in \{\pm 1\}^k} \prod_{\sigma \in \Delta_k} \frac{1}{\mathcal{V}_\sigma} \mathcal{O}_\Gamma(l_e, l_{e'}^\epsilon(l_e)) \quad (10)$$

where we have dropped out the overall factor which corresponds to the gauge volume.

Let us conclude this part: we have started with a Feynman amplitude on flat space-time; in the usual formulation, information about flat geometry is encoded in the Lebesgue measure. By working with the invariant measure, we have moved this information in the weight assigned to the triangulation, and in the relations $l_{e'}^\epsilon(l_e)$ between ‘missing’ lengths, labels and orientations. Our goal now is to show that this information about flat geometry can be encoded entirely into a dynamical topological state sum (spin foam) model.

III. FEYNMAN DIAGRAMS AS SPIN FOAM MODELS

A. The main result

One of the main results of this paper is the fact that the Feynman amplitude (10) can be written as the expectation value of an observable

$$I_\Gamma = \langle \mathcal{O}_\Gamma(l_e) \rangle_{Z_\Delta}, \quad \mathcal{O}_\Gamma = \prod_{e \in \Gamma} G^F(l_e) \quad (11)$$

for the spin foam model:

$$Z_\Delta = \frac{1}{(2\pi)^{|e|}} \int \prod_{e \in \Delta} d l_e l_e^2 \sum_{\{s_e, \epsilon_\tau\}} \left(\prod_{\tau} \frac{e^{i\epsilon_\tau S_\tau(s_e, l_e)}}{\mathcal{V}_\tau} \right) \quad (12)$$

Δ is an arbitrary triangulation of a closed 3D-manifold and Γ is embedded into the one-skeleton of Δ . $|e|$ is the number of edges of Δ . The edges carry a double label (l_e, s_e) , where l_e is a real positive number and s_e is an integer. These labels are summed over a domain where triangular inequalities are satisfied for the l_e ’s. $\epsilon_\tau = \pm 1$ denotes the orientation of the tetrahedron τ . The product in parenthesis is over all tetrahedra. The action for each tetrahedron is similar to the Regge action [16]

$$S_\tau(s_e, l_e) = \sum_{e \in \tau} s_e \theta_e^\tau(l_e), \quad (13)$$

with $\theta_e^\tau(l_e)$ the interior dihedral angle of the edge e in τ . The observable is simply the product of propagators in which the distances are replaced by the labels l_e living on the graph Γ . The equality

(11) is obtained if one restricts to trivial topologies (Δ is a triangulation of the 3-sphere S^3). Note that all edges are summed over, and therefore no reference is made to flat geometry.

Furthermore, we will show that this model is topological and that the weight associated to it can be interpreted algebraically as the $6j$ -symbol of the Poincaré group. The rest of the paper is devoted to the demonstration of these statements. The notations we use are summarized and explained in details in appendix A.

B. Emergence of the model

1. A key identity

The keystone of the result announced above is a geometric identity associated with the possibility to embed a 4-simplex in $\mathbb{R}^3$. A similar identity appears in [14, 17] as a basis to construct 3D manifold invariants. Let $G \equiv \mathcal{V}^2$ be the square of the volume of a 4-simplex $(0, \dots, 4)$ with edge lengths l_{ij} . We denote by τ_j the tetrahedron obtained by dropping the point j of the 4-simplex, and by $\mathcal{V}_j$ its volume. We will consider both the complex of the three tetrahedra τ_1, τ_2 and τ_3 surrounding the edge (04) , and the complex of the two tetrahedra τ_0, τ_4 sharing the face $[123]$ (see Fig 2). The orientation of τ_j is denoted by ϵ_j .

We work with fixed lengths $l_{ij}, (ij) \neq (04)$, the length l_{04} being free to fluctuate. Then, the following identity of measures holds:

$$4l_{04}\delta(G) = \sum_{\epsilon_0, \epsilon_4} \frac{\delta(l_{04} - l_{04}^{\epsilon_0\epsilon_4})}{\mathcal{V}_0 \mathcal{V}_4} = \sum_{\epsilon_1, \epsilon_2, \epsilon_3} l_{04}^2 \frac{\delta(\omega_{04}^\epsilon(l_{04}))}{\mathcal{V}_1 \mathcal{V}_2 \mathcal{V}_3}. \quad (14)$$

The delta function in the third term is the 2π -periodic delta function. The quantity ω_{04}^ϵ , considered as a function of l_{04} and orientations, is the deficit angle carried by the edge (04) , defined to be:

$$\omega_{04}^\epsilon = \sum_{j=1}^3 \epsilon_j \theta_{04}^j$$

where $\theta_{04}^j \in [0, \pi]$ is the interior dihedral angle of the edge (04) in τ_j . It represents the curvature localized on the edge, in the sense of Regge calculus. It vanishes modulo⁴ 2π if and only if the complex of three tetrahedra (τ_1, τ_2, τ_3) can be mapped in $\mathbb{R}^3$ with the orientations ϵ_j . Thus, the delta function of the deficit angle acts as a projector on the space of flat geometries.

On the other hand, the complex of the two tetrahedra τ_0, τ_4 can always be mapped in $\mathbb{R}^3$, in a unique (modulo translations and rotations) way depending on the orientations. The ‘missing’ distance between the points 0,4 is then a well defined function $l_{04}^{\epsilon_0\epsilon_4}$ of l_{ij} and the product orientations.

We now give a proof of (14). We will need three intermediate results: the first one states that the equation

$$G(l_{04}) = 0,$$

⁴ If we don’t work modulo 2π the condition for the existence of an embedding of the 4-simplex $(0, \dots, 4)$ in flat space is $\omega_{04}^\epsilon = 2\pi n(\epsilon)$ where $n(\epsilon) \in \{0, +1, -1\}$ depends on the orientations in this way:

$$n(\epsilon) = \begin{cases} \epsilon_1 & \text{if } \epsilon_i = \epsilon_j \quad \forall (i, j) \\ 0 & \text{if not} \end{cases} \quad (15)$$




FIG. 2: Pentagonal representation of a complex of two tetrahedra 0, 4 sharing a face [123], and a complex of three tetrahedra 1, 2, 3 sharing an edge (04).

where $G \equiv \mathcal{V}^2$ is considered to be a function of l_{04} , all the other lengths being fixed, admits exactly two solutions $l_{04}^\pm$. It was implicit in our discussion above, where we have mentioned, moreover, that these solutions are labelled precisely by the product $\epsilon_0 \epsilon_4$. This result is geometrically clear since for each value of l_{04} one can embed the simplex in $\mathbb{R}^4$ in such a way to map, say, τ_0 into a fixed tetrahedron. The volume of the simplex vanishes only when the vertex 0 belongs to the space E spanned by the tetrahedron τ_0 . As l_{04} varies within its domain, the vertex 0 moves along a circle around the face [123], intersecting E at exactly two points.

We can also give an algebraic justification for having only two solutions. Indeed, G is the Cayley-Menger determinant of the 4-simplex. The Cayley-Menger matrix is a 6×6 symmetric matrix such that its diagonal elements are 0 and the off diagonal elements are given by $L_{ab} \equiv -l_{ab}^2/2$, except for the first column and row where they are 1. Its determinant is therefore a quadratic function in L_{04} . Moreover, the coefficients of this polynomial can be computed explicitly (see (A13) and (A15)):

$$\frac{\partial G}{\partial L_{04}} = -2\mathcal{V}_0\mathcal{V}_4 \cos \theta_{04}, \quad \frac{\partial^2 G}{\partial L_{04}^2} = -2\mathcal{V}_{04}^2, \quad (16)$$

where θ_{04} is the interior dihedral angle between the two tetrahedra 0 and 4, and $\mathcal{V}_{04}$ is the area of the face [123].

The second result states that the equation

$$\omega_{04}^\epsilon(l_{04}) = 0,$$

where ω is considered as a function of l_{04} (the other lengths being fixed) and orientations $\epsilon_1, \epsilon_2, \epsilon_3$, admits exactly four solutions. Indeed this equation expresses the condition of existence of a map of the 4-simplex in $\mathbb{R}^3$ giving orientations $\epsilon = \epsilon_1, \epsilon_2, \epsilon_3$ to the corresponding tetrahedra. Hence, first, it imposes $G = 0$, and therefore, according to the previous result $l_{04} = l_{04}^\pm$. Now the data $\{l_{ij}, l_{04}^\pm\}$ determines the geometry of the complex in $\mathbb{R}^3$ modulo orientation reversal in this space (in other words simultaneous change of all orientations). Therefore for each $l_{04}^\pm$, there exist exactly two sets of orientations $\eta \epsilon_i^\pm$, $\eta \in \{\pm 1\}$ for which the equation is satisfied, which proves the statement.

Let $(l_{04}^\pm, \epsilon_i)$ denote one of the four solutions of $G = \omega = 0$. Then a variation δl_{04} of the length induces variations of δG of the volume and $\delta \omega$ of the deficit angle. Our third lemma, established

in appendix A, relates these variations to each other:

$$\left. \frac{\partial G}{\partial \omega_{04}^\epsilon} \right|_{l_{04}^\pm} = -2 \epsilon_1 \epsilon_2 \epsilon_3 \frac{\mathcal{V}_1 \mathcal{V}_2 \mathcal{V}_3}{l_{04}} \quad (17)$$

$$\left. \frac{\partial G}{\partial l_{04}} \right|_{l_{04}^\pm} = \mp 2 \mathcal{V}_0 \mathcal{V}_4 l_{04} \quad (18)$$

The identity (14) is obtained by evaluating in two different ways the functional $\delta(G)$. We first consider G to be a function of l_{04} . We have:

$$\delta(G) = \sum_{\epsilon \in \{\pm 1\}} \delta(l_{04} - l_{04}^\epsilon) \left| \frac{\partial G}{\partial l_{04}} \right|^{-1} = \sum_{\epsilon_0, \epsilon_4} \frac{1}{4l_{04}} \frac{1}{\mathcal{V}_0 \mathcal{V}_4} \delta(l_{04} - l_{04}^{\epsilon_0 \epsilon_4}). \quad (19)$$

In order to get the second equality we have used (18), replaced the sum over the solutions by a sum over the orientations of the tetrahedra 0, 4, taking into account the overcounting by a factor 2.

Instead of considering G as a function of l_{04} , one can consider it to be a function which depends on the orientations $\epsilon_1, \epsilon_2, \epsilon_3$ and the corresponding deficit angle ω_{04}^ϵ . The following property

$$G = 0 \iff \exists \epsilon_1, \epsilon_2, \epsilon_3 \text{ such that } \omega_{04}^\epsilon = 0 \pmod{2\pi}.$$

yields:

$$\delta(G) = \frac{1}{2} \sum_{\epsilon_1, \epsilon_2, \epsilon_3} \delta(\omega_{04}^\epsilon) \left| \frac{\partial G}{\partial \omega_{04}^\epsilon} \right|^{-1} = \sum_{\epsilon_1, \epsilon_2, \epsilon_3} \frac{l_{04}}{4} \frac{\delta(\omega_{04}^\epsilon)}{\mathcal{V}_1 \mathcal{V}_2 \mathcal{V}_3} \quad (20)$$

The factor 1/2 in the first equality comes from the fact that, if $\omega_{04}^\epsilon = 0$, then $\omega_{04}^{-\epsilon} = 0$, so when we sum over all ϵ_i we are overcounting with a factor of 2. The equality between (19) and (20) leads to the identity (14).

It is worth noting that one can actually derive an identity slightly stronger than (14). In fact, (18) divided by (17) gives the derivative of the deficit angle with respect to the length:

$$\left. \frac{\partial \omega_{04}^\epsilon}{\partial l_{04}} \right|_{l_{04}^\pm} = \pm \epsilon_1 \epsilon_2 \epsilon_3 l_{04}^2 \frac{\mathcal{V}_0 \mathcal{V}_4}{\mathcal{V}_1 \mathcal{V}_2 \mathcal{V}_3} \quad (21)$$

It is then straightforward to check that

$$\forall \eta, \quad \frac{f(l_{04}^\pm)}{\mathcal{V}_0 \mathcal{V}_4} = \int_{l_{04}^\pm} dl_{04} l_{04}^2 \frac{\delta(l_{04} - l_{04}^\pm)}{\mathcal{V}_1 \mathcal{V}_2 \mathcal{V}_3} \left| \frac{\partial \omega_{04}^{\eta^\pm}}{\partial l_{04}} \right|^{-1} f(l_{04}) \quad (22)$$

$$= \int_{l_{04}^\pm} dl_{04} l_{04}^2 \frac{\delta(\omega_{04}^{\eta^\pm})}{\mathcal{V}_1 \mathcal{V}_2 \mathcal{V}_3} f(l_{04}). \quad (23)$$

where the label appearing as an index of integration means that one integrates in the neighborhood of the value $l_{04}^\pm$.

In the next two parts of this section, we will take advantage of this identity to express the Feynman graph amplitude as a spin foam amplitude.

2. Spin foam in Feynman graph: inducing the geometry dynamically

In order to understand the mechanism, let us begin by studying a simple example. Suppose that our Feynman graph Γ has the structure depicted in Fig 2: it admits five vertices and ten links. The two tetrahedra τ_0, τ_4 form a triangulation of a 3-ball. According to the construction described in section II, the amplitude of this graph is:

$$I_\Gamma = \int \prod_{(ij) \neq (04)} l_{ij} dl_{ij} \sum_{\epsilon_0, \epsilon_4} \frac{1}{\mathcal{V}_0 \mathcal{V}_4} \mathcal{O}_\Gamma(l_{ij}, l_{04}^{\epsilon_0 \epsilon_4}(l_{ij})) \quad (24)$$

where $\mathcal{O}_\Gamma$ is the product of propagators

$$G^F(l_{04}^{\epsilon_0 \epsilon_4}(l_{ij})) \prod_{(ij) \neq (04)} G^F(l_{ij})$$

The length of (04) depends explicitly on the nine other lengths. The previous identity (14) allows us to remove this dependence by introducing a new label l_{04} and summing over it. Plugging it into (24), and expanding the periodic delta function, as a sum $2\pi\delta(\omega) = \sum_{s \in \mathbb{Z}} e^{is\omega}$, we get:

$$I_\Gamma = \frac{1}{2\pi} \int \prod_{(ij) \neq (04)} l_{ij} dl_{ij} \int dl_{04} l_{04}^2 \sum_{s_{04} \in \mathbb{Z}} \sum_{\epsilon_1, \epsilon_2, \epsilon_3} \frac{e^{is_{04}\omega_{04}^\epsilon}}{\mathcal{V}_1 \mathcal{V}_2 \mathcal{V}_3} \mathcal{O}_\Gamma(l_{ij}, l_{04}) \quad (25)$$

Notice that the tetrahedra τ_1, τ_2, τ_3 form a new triangulation of the ball. In this expression, one begins to recognize the structure of the model (14), since the total action can be expressed in terms of the deficit angles associated with the edges of Δ

$$S_\Delta \equiv \sum_\tau \epsilon_\tau S_\tau = \sum_e s_e \omega_e^\epsilon. \quad (26)$$

Let us summarize where we stand so far in addressing the first step, which was to transfer the information about flat geometry from the measure into relations $l_e^\epsilon(l_e)$; it has led us from (1) to (10). Now considering two tetrahedra sharing a face in the triangulation Δ_k , we have learned from the example above that the ‘missing’ distance variable can be replaced by a free label that we integrate over. The price to pay is the apparition of a constraint term - the delta function of the deficit angle - which, in the language of Regge calculus, imposes locally, the geometry to be flat. Ultimately, by expanding the delta function, we begin to see a model emerging from the Feynman graph, where the flat geometry is induced dynamically.

The result (11) will be proved in two steps. First, we will interpret our key identity as expressing the invariance under Pachner moves, and thus establish the topological invariance of the model (12). Secondly we will write the measure of the Feynman graph as the partition function of this model, computed on a specific triangulation.

C. Topological invariance

In this section we show that the spin foam model (12),

$$Z_\Delta = \frac{1}{(2\pi)^{|e|}} \int_{\text{GF}} \prod_{e \in \Delta} dl_e l_e^2 \sum_{\{s_e, \epsilon_\tau\}} \left(\prod_\tau \frac{e^{is_\tau S_\tau(s_e, l_e)}}{\mathcal{V}_\tau} \right), \quad (27)$$

is independent of the choice of the triangulation and depends only on the topology of the manifold that Δ triangulates. The label (GF) appearing as an index of integration means that the integrals have to be *gauge fixed*, in order to properly define the partition function.

The necessity of gauge fixing the model is manifest if one carefully looks at the integrand of (12). In fact, the sum over integers s of the exponentials leads to a product over edges of delta functions $\delta(\omega_e^\epsilon)$. The constraints $\omega_e^\epsilon = 0 \mod 2\pi$ are not all independent however, and the redundancy of delta functions induces divergences. These naive divergences are related to the existence of gauge symmetries in the model, which have to be gauge fixed. This is analogous to what happens in the study of the 3d quantum gravity amplitudes, as explained in [18]. The symmetries and the corresponding gauge fixing of the model will be studied in great detail in the next section .

In order to specify this gauge fixing, we first choose 4 vertices which belong to a tetrahedron τ_0 of Δ , and then assign to any other internal vertex v of Δ a tetrahedron τ_v to which this vertex belongs. This assignment is said to be admissible if the following condition is met: given two vertices v_1 and v_2 of Δ that are not in τ_0 , then $(v_1 v_2) \not\subset \tau_{v_1} \cap \tau_{v_2}$, that is, the edge $(v_1 v_2)$ (when it exists) does not belong to the intersection of the corresponding tetrahedra. We denote such an admissible gauge fixing assignment $(\tau_0, \{\tau_v\}_{v \notin \tau_0})$ by T . We say that a tetrahedron τ belongs to T if either $\tau = \tau_0$ or $\tau = \tau_v$ for some vertex not in τ_0 ; we say that an edge e belongs to T if either $e \in \tau_0$ or $e \in \tau_v$, and if e admits v as one of its vertex - for each τ_v there are three such edges, carrying the labels l_{v1}, l_{v2}, l_{v3} . The admissibility condition insures that these triplets of edges are independent, that is, we cannot have $l_{vi} = l_{v'j}$. We will see later that the admissibility requirement can always be met.

We now define a gauge fixing term δ_{GF}^T and a Faddeev-Popov determinant D_{FP}^T as

$$\delta_{GF}^T(l_e, s_e) = \prod_{e \in T} \delta(l_e - l_e^o) (2\pi) \delta_{s_e, s_e^o} \quad (28)$$

$$D_{FP}^T(l_e, s_e) = \prod_{\tau \in T} \frac{1}{2} \left(\frac{\mathcal{V}_\tau}{l_{v1} l_{v2} l_{v3}} \right)^2. \quad (29)$$

The gauge fixing term imposes the values of l_e and s_e to be l_e^o, s_e^o , which are arbitrary fixed. Fixing the labels s 's living on the edges of the assignment removes the redundant delta functions of the deficit angle at these edges. Now note that the constraints $\omega_e^\epsilon(l) = 0 [2\pi]$ act both on the labels l 's and the orientations. Therefore by removing them, we not only remove redundant constraints on the l 's but also necessary restrictions acting on orientations. As shown in appendix, one can reintroduce these restrictions by inserting an additional gauge fixing factor⁵ in δ_{GF}^T :

$$\prod_{v \in T} \chi(\Omega_v^\epsilon(l_e)).$$

The function $\chi(x)$ is a characteristic function defined to be constant, with value 1, on $4\pi\mathbb{Z}$, and 0 elsewhere; and $\Omega_v^\epsilon = \sum_{\tau \supset v} \epsilon_\tau A_\tau$ is the algebraic sum, over all tetrahedra sharing the vertex v , of the solid angle A_τ at v within τ .

The gauge fixed partition function is obtained by inserting in the integral (27) the product $\delta_{GF}^T D_{FP}^T$.

Showing the topological invariance of the model amounts to showing the invariance of Z_Δ under the Pachner moves (2, 3) and (1, 4), and its independence of the choice of T . The invariance under

⁵ This factor acts ‘on-shell’ as Kronecker symbols for the orientations and is understood here as a gauge fixing. An interesting open question is whether this constraint can be expressed off-shell and implemented dynamically.

the move (2, 3) follows from the pentagonal identity,

$$\sum_{\epsilon_0, \epsilon_4} \frac{e^{i\epsilon_0 S_0}}{\mathcal{V}_0} \frac{e^{i\epsilon_4 S_4}}{\mathcal{V}_4} = \frac{1}{2\pi} \sum_{\epsilon_1, \epsilon_2, \epsilon_3} \int dl_{04} l_{04}^2 \sum_{s_{04}} \frac{e^{i\epsilon_1 S_1}}{\mathcal{V}_1} \frac{e^{i\epsilon_2 S_2}}{\mathcal{V}_2} \frac{e^{i\epsilon_3 S_3}}{\mathcal{V}_3}, \quad (30)$$

where $\tau_0, \dots, \tau_4$ labels the five tetrahedra which triangulate the boundary of a 4-simplex, τ_i being the tetrahedron where the vertex i is omitted; the invariance under the move (1, 4) is due to the (1, 4) identity

$$\sum_{\epsilon_0} \frac{e^{i\epsilon_0 S_0}}{\mathcal{V}_0} = \frac{1}{(2\pi)^4} \sum_{\epsilon_j} \int \prod_{j=1}^4 dl_{0j} l_{0j}^2 \sum_{s_{0j}} \prod_{j=1}^4 \frac{e^{i\epsilon_j S_j}}{\mathcal{V}_j} \delta_{\text{GF}} D_{\text{FP}} \quad (31)$$

δ_{GF} denotes the gauge fixing term and D_{FP} the Faddeev-Popov determinant:

$$\delta_{\text{GF}} = (2\pi)^3 \prod_{i=1}^3 \delta(l_{0i} - l_{0i}^o) \delta_{s_{0i}, s_{0i}^o} \chi(\Omega_0^\epsilon), \quad D_{\text{FP}} = \frac{\mathcal{V}_4^2}{2 \prod_{i=1}^3 l_{0i}^2} \quad (32)$$

where l_{0i}^o, s_{0i}^o are any fixed values. At this stage the insertion of these terms are needed for topological invariance; they will be fully and independently justified in the next section (together with the gauge fixing denomination).

The proof of (30) is a simple adaptation of the argument leading to the key identity (14). As before, we denote by $\eta\epsilon_i^\pm, \eta = \pm 1$, the two sets of orientations for which the deficit angle $\omega_{04}^\epsilon(l_{04}^\pm)$ vanishes. If this dihedral angle is zero modulo 2π , the 4-simplex [01234] can be mapped in $\mathbb{R}^3$. Each edge (ab) belongs then to a complex of three tetrahedra embedded in $\mathbb{R}^3$. One can therefore find orientations $\epsilon_0^\pm, \epsilon_4^\pm$ of τ_0, τ_4 , for which all deficit angles $\omega_{ab}^{\epsilon^\pm}$ are also zero modulo 2π . The actions of the (2, 3) move evaluated for $l_{04} = l_{04}^\pm$ are then related by

$$\epsilon_0^\pm S_0 + \epsilon_4^\pm S_4 + \epsilon_1^\pm S_1^o + \epsilon_2^\pm S_2^o + \epsilon_3^\pm S_3^o \equiv \sum_{(ab) \neq (04)} s_{ab} \omega_{ab}^{\epsilon^\pm} = 0 \quad [2\pi] \quad (33)$$

The superscript o means that $s_{04} = 0$ in the actions. This equality, together with (23), show that

$$\forall \eta, \quad \frac{e^{i\eta\epsilon_0^\pm S_0}}{\mathcal{V}_0} \frac{e^{i\eta\epsilon_4^\pm S_4}}{\mathcal{V}_4} = \int_{l_{04}^\pm} dl_{04} l_{04}^2 \delta(\omega_{04}^{\eta\epsilon^\pm}) \prod_{i=1}^3 \frac{e^{-i\eta\epsilon_i^\pm S_i^o}}{\mathcal{V}_i} \quad (34)$$

Now we want to sum over $\eta = \pm 1$. Since $\eta\epsilon_i^\pm$ are the only possible choice of orientations for which $\omega_{04}^\epsilon = 0$ admits a solution around $l_{04}^\pm$, one can replace, in the RHS, the summation over η by a sum over all orientations. Then summing the contributions of the two values $l_{04}^\pm$ leads to

$$\sum_{\pm} \sum_{\eta} \frac{e^{i\eta\epsilon_0^\pm S_0}}{\mathcal{V}_0} \frac{e^{i\eta\epsilon_4^\pm S_4}}{\mathcal{V}_4} = \sum_{\epsilon_1, \epsilon_2, \epsilon_3} \int dl_{04} l_{04}^2 \delta(\omega_{04}^\epsilon) \prod_{i=1}^3 \frac{e^{i\epsilon_i S_i^o}}{\mathcal{V}_i}. \quad (35)$$

Now it is easy to check⁶ that the summation in the LHS can be replaced by a sum over the values of orientations ϵ_0, ϵ_4 . If one now expands the delta function in the RHS as a sum over s_{04} of $1/2\pi \exp(is_{04}\omega_{04})$, one obtains (30). It's worth noting that we could have inserted any function $f(l_{04})$ of the length l_{04} in both sides of all our equalities, without modification, and therefore (30)

⁶ In fact, our definitions of orientations (see appendix) imply that $\epsilon_0^\pm \epsilon_4^\pm = \pm 1$. It shows that the application $(\pm, \eta) \mapsto (\eta\epsilon_0^\pm, \eta\epsilon_4^\pm)$ is surjective onto $\{-1, +1\}$.

should be understood as an equality of measures on the space of such functions. Since S_4 and $\mathcal{V}_4$ do not depend on l_{04} , we multiply both sides of (34) by $\mathcal{V}_4 e^{-\eta \epsilon_4^\pm S_4}$, to get

$$\forall \eta, \quad \frac{e^{\eta \epsilon_0^\pm S_0}}{\mathcal{V}_0} = \int_{l_{04}^\pm} dl_{04} l_{04}^2 \delta(\omega_{04}^{\eta \epsilon^\pm}) \prod_{i=1}^3 \frac{e^{-\eta \epsilon_i^\pm S_i^\circ}}{\mathcal{V}_i} \frac{e^{-\eta \epsilon_4^\pm S_4}}{\mathcal{V}_4} \mathcal{V}_4^2, \quad (36)$$

where in the RHS the values of $l_{4i} \equiv l_{4i}^\circ$ are fixed to be any value - since the reasoning leading to (34) does not depend on it. We now insert the trivial ‘gauge fixing’ identities

$$1 = \int \prod_{i=1}^3 dl_{0i} l_{0i}^2 \prod_{i=1}^3 \frac{\delta(l_{0i} - l_{0i}^\circ)}{l_{0i}^2} \quad (37)$$

$$1 = \sum_{\epsilon_4} \frac{1}{(2\pi)^3} \sum_{s_{0i}} \prod_{i=1}^3 \delta_{s_{0i}, s_{0i}^\circ} (2\pi)^3 \chi(\Omega_v^\epsilon), \quad (38)$$

where $\epsilon = (\eta \epsilon_i^\pm, \epsilon_4)$. As can be easily seen geometrically, and as more precisely shown in appendix, the algebraic solid angle Ω_0^ϵ , evaluated at $l_{04}^\pm$, vanishes (modulo 4π) only if $\epsilon_4 = \eta \epsilon_4^\pm$, and therefore its image by the function χ acts as a Kronecker symbol for ϵ_4 . Hence, we obtain

$$\frac{1}{2} \frac{e^{\eta \epsilon_0^\pm S_0}}{\mathcal{V}_0} = \frac{1}{(2\pi)^4} \sum_{\epsilon_4} \int \prod_{i=1}^3 dl_{0i} l_{0i}^2 \int_{l_{04}^\pm} dl_{04} l_{04}^2 \sum_{s_{0i}} \delta(\omega_{04}^{\eta \epsilon^\pm}) \prod_{i=1}^3 \frac{e^{-\eta \epsilon_i^\pm S_i^\circ}}{\mathcal{V}_i} \frac{e^{-\eta \epsilon_4^\pm S_4}}{\mathcal{V}_4} \delta_{GF} D_{FP}. \quad (39)$$

for all $\eta = \pm 1$. Eventually, summing as before both sides of this equality over η , adding the contributions of l_{04}^+ and l_{04}^- , and expanding the delta function in the RHS, show (31).

D. Symmetries and gauge fixing

We have seen in the previous part that, in order to have a model which is topologically invariant, we need to insert gauge fixing and Faddeev-Popov terms in the definition of the spin foam model (12). We now want to show that these factors are not only needed for topological invariance, but arise naturally from the gauge fixing of the symmetries of the action.

The action that arises from our construction reads:

$$S_\Delta[s_e, l_e] = \sum_\tau \epsilon_\tau S_\tau(s_e, l_e) = \sum_e s_e \omega_e^\epsilon(l_e) \quad (40)$$

The naive divergences of the model (12) can be seen as arising from symmetries of the model mapping solutions to solutions. Fixing these symmetries, treated as gauge symmetries, will lead to the Faddeev-Popov determinants described in the previous section. In order to characterize these symmetries infinitesimally, we will study the zero modes of the Hessian of S_Δ . Here s_e will be treated as a continuous variable, in order to simplify the analysis⁷.

Let us first describe the classical solutions. The equations of motion (eom) are

$$0 = \frac{\delta S}{\delta s_e} = \omega_e^\epsilon(l) \quad \forall e \quad (41)$$

$$0 = \frac{\delta S}{\delta l_e} = \sum_\tau \epsilon_\tau \left(\sum_{e' \in \tau} s_{e'} \frac{\delta \theta_{e'}^\tau}{\delta l_e} \right) = \sum_{e'} s_{e'} \frac{\delta \omega_{e'}^\epsilon}{\delta l_e} \quad \forall e \quad (42)$$

⁷ We can treat s_e as a continuous variable if one authorizes ω_e to be shifted by $2n\pi$ for n an arbitrary integer. This is clear from the expression of the periodic delta function $\delta(\omega) = \sum_n \int ds e^{is(\omega - 2n\pi)}$. This affects the classical solutions but not the equations that involve derivatives of ω , which are our main interest here.

The first equation expresses the flatness condition. A set of lengths $\{l_e^0\}$ is solution if the complex can be locally⁸ mapped in $\mathbb{R}^3$ in a way that respects orientations ϵ . The second equation is reminiscent of the *Schläfli identity* for flat tetrahedra

$$\sum_{e' \in \tau} l_{e'} \frac{\delta \theta_{e'}^\tau}{\delta l_e} = 0 \rightarrow \sum_{e'} l_{e'} \frac{\delta \omega_{e'}^\epsilon}{\delta l_e} = 0. \quad (43)$$

This shows that a solution of (42) is given by the values $s_e^0 = \alpha l_e$, with α an arbitrary constant. Notice that if one inserts this solution into the action (40), one recovers the Regge action of discrete 3D gravity

$$\mathcal{S}_R = \alpha \sum_e l_e \omega_e^\epsilon(l_e) \quad (44)$$

In the following a solution $\{l_e^0, s_e^0 = l_e^0\}$ of the eom will be called a Regge solution (we put the overall scale factor $\alpha = 1$ for simplicity).

Zeros modes are infinitesimal transformations of the fields (here the labels) that belong to the kernel of the Hessian $\delta^2 S$, computed on shell. This kernel is characterized by deformations $\delta s_e, \delta l_e$ that satisfy the equations

$$\sum_{e'} \frac{\delta \omega_e}{\delta l_{e'}} \delta l_{e'} = 0 \quad \forall e \quad (45)$$

$$\sum_{e', e''} s_{e''} \frac{\delta^2 \omega_{e''}}{\delta l_e \delta l_{e'}} \delta l_{e'} + \sum_{e'} \frac{\delta \omega_{e'}}{\delta l_e} \delta s_{e'} = 0 \quad \forall e \quad (46)$$

where the label ϵ has been dropped for clarity. This system of equations can be rearranged if one uses the derivative of the Schläfli identity

$$\frac{\delta}{\delta l_e} \left[\sum_{e''} l_{e''} \frac{\delta \omega_{e''}}{\delta l_{e'}} \right] = \frac{\delta \omega_e}{\delta l_{e'}} + \sum_{e''} l_{e''} \frac{\delta^2 \omega_{e''}}{\delta l_e \delta l_{e'}} = 0, \quad (47)$$

and the fact that the matrix $\left(\frac{\delta \omega_e}{\delta l_{e'}} \right)$ is symmetric, as the Hessian of the Regge fonction (44). We can write (46) as

$$\sum_{e', e''} (s_{e''} - l_{e''}) \frac{\delta^2 \omega_{e''}}{\delta l_e \delta l_{e'}} \delta l_{e'} + \sum_{e'} \frac{\delta \omega_e}{\delta l_{e'}} \delta (s_{e'} - l_{e'}) = 0. \quad (48)$$

If one focuses on fluctuations around Regge solutions, the system becomes equivalent to

$$\sum_{e'} \frac{\delta \omega_e}{\delta l_{e'}} \delta l_{e'} = \sum_{e'} \frac{\delta \omega_e}{\delta l_{e'}} \delta s_{e'} = 0 \quad \forall e \quad (49)$$

One can easily find infinitesimal variations $\delta l_e, \delta s_e$ solutions of (49). Indeed, an embedding in $\mathbb{R}^3$ of the complex Δ , with the lengths $\{l_e^0\}$, gives positions to the vertices. Infinitesimal moves of the vertices in $\mathbb{R}^3$ around these positions induces deformations of the edge lengths. By construction, the new deficit angles still vanish. Thus, these deformations correspond to one of the symmetries

⁸ A global map exists, and is unique modulo translations and rotations, if the manifold that Δ triangulates is simply connected [17].

we are considering, acting on vertices of the triangulation. The positions of the vertices in an embedding are therefore pure gauge in our model. The generators are 3-vectors $\vec{\alpha}_v$ attached to each vertex v which induce displacements of v in $\mathbb{R}^3$. They determine variations $\delta s_e = \delta l_e(\vec{\alpha}_v, l_e^0)$, $\delta l_e = \delta l_e(\vec{\beta}_v, l_e^0)$, which are solutions to the system of equations (49).

These transformations map solutions to solutions⁹; the corresponding symmetry group is $(\mathbb{R}^3 \times \mathbb{R}^3)^{|v|}$, where $|v|$ is the number of vertices of the triangulation. The gauge fixing is performed by fixing a subset of edge lengths while taking into account the Jacobian or Faddeev-Popov determinant. These Jacobians can easily be read out from the results of section II. We start by choosing four vertices that form a tetrahedron τ_0 in the triangulation. For these vertices we have

$$d^3\alpha_1 d^3\alpha_2 d^3\alpha_3 d^3\alpha_4 = 2 d^3\alpha d\Lambda \frac{\prod_{i<j} l_{ij} dl_{ij}}{\mathcal{V}_{\tau_0}}. \quad (50)$$

The factor 2 is due to the sum over values of the orientation for τ_0 . For each additional vertex v , we choose a tetrahedron τ_v , and denote $l_{v_1}, l_{v_2}, l_{v_3}$ the three edges of τ_v touching v . In this case we have

$$d^3\alpha_v = 2 \frac{\prod_i l_{v_i} dl_{v_i}}{\mathcal{V}_{\tau_v}}. \quad (51)$$

Each time we add a new vertex v_n , one should insure that τ_{v_n} is picked in such a way that the three corresponding edges do not overlap with any of the previously chosen edges $l_{v_{1i}}, \dots, l_{v_{n-1i}}$, in order for an edge to be selected at most once in this process. We are going to see in the following how this can always be concretely implemented for any closed triangulation.

This procedure specifies an admissible gauge fixing assignment T_l , as defined in the previous section, which allows us to gauge fix the symmetry acting on l 's. The gauge fixing term is given by

$$\delta_{GF}^{T_l} = \prod_{e \in T_l} \delta(l_e - l_e^o) \quad (52)$$

and the determinant by

$$(D_{FP}^{T_l})^{-1} \equiv \frac{\bigwedge_v d^3\alpha_v}{d^3\alpha d\Lambda \bigwedge_{e \in T_l} dl_e} = 2^{|v|-3} \frac{\prod_{e \in T_l} l_e}{\prod_{\tau \in T_l} \mathcal{V}_{\tau}}, \quad (53)$$

where $|v|$ is the number of vertices of Δ . The same procedure is applied to the gauge fixing of the symmetry acting on s 's, with a gauge fixing assignment T_s , and the corresponding gauge fixing term and determinant

$$\delta_{GF}^{T_s} = \prod_{e \in T_s} (2\pi) \delta_{s_e, s_e^o} \prod_{v \in T_s} \chi(\Omega_v^\epsilon(l_e)) \quad (54)$$

$$(D_{FP}^{T_s})^{-1} \equiv \frac{\bigwedge_v d^3\alpha_v}{d^3\alpha d\Lambda \bigwedge_{e \in T_s} ds_e} = \frac{\prod_{e \in T_s} l_e}{\prod_{\tau \in T_s} \mathcal{V}_{\tau}}. \quad (55)$$

There is no factor 2 in this Faddeev-Popov determinant, since here orientations of the s 's tetrahedra are fixed thanks to the factors $\chi(\Omega)$ in the gauge fixing term. The insertion of the functions χ

⁹ These symmetries, acting on the space of solutions, do not however leave the full action invariant away from classical configurations. Whether or not these symmetries admit an off-shell extension is a interesting open question not addressed here.

restricts the values of orientations to those for which solutions of the eom (41), around which the zeros modes which have to be fixed, exist. One thus recovers the gauge fixing described in the previous section when one chooses the same gauge fixing assignment $T_l = T_s$ for l and s .

Finally, we need to show that an admissible gauge fixing assignment can always be reached. One way to construct such an assignment is to choose a rooted tree (also denoted T for simplicity) in the one skeleton of the dual triangulation. The one skeleton of the dual triangulation is a 4-valent graph; each vertex of this graph corresponds to a tetrahedron of Δ . A tree is a connected subgraph of this skeleton which contains no loop. A root is a choice of one particular end point of this tree diagram. We use the notation v^* for the vertices of the dual triangulation, and τ_{v^*} for the tetrahedron of Δ dual to v^* . A tree T is said *maximal* if each vertex of Δ belongs to a tetrahedron dual to a vertex of T : $\forall v \in \Delta, \exists v^* \in T, v \subset \tau_{v^*}$.

Given a maximal rooted tree T one can construct an admissible assignment as follows. The root of T corresponds to a choice of an initial tetrahedron τ_0 . Then, lets pick an edge e^* that goes from the root to a nearby tetrahedron τ_{e^*} : this edge is dual to a face f_{e^*} of the triangulation. If we consider $\tau_{e^*} - f_{e^*}$, this consists of three edges meeting at a vertex $v_{e^*} \in \tau_{e^*}$; in this way we define a correspondence between e^* and a pair (τ_{e^*}, v_{e^*}) . This correspondence can be extended to the all tree if one uses the natural orientation of the rooted tree. Indeed, since the tree is rooted, every edge of the tree can be oriented with the orientation always going farther away from the root. And to any edge e^* of T we assign a tetrahedron τ_{e^*} corresponding to the end point of this edge and a vertex v_{e^*} being the unique vertex in $\tau_{e^*} - f_{e^*}$. This assignment is clearly admissible. In order to see why, lets suppose that there exists an edge of Δ that belongs to both $\tau_{e^*} - f_{e^*}$ and $\tau_{e'^*} - f_{e'^*}$. This edge would belong to the intersection; but this intersection is empty since tetrahedra always intersect on faces dual to the rooted tree. Eventually, the maximality of the tree implies the maximality of the gauge fixing assignment.

E. Observables and partial gauge fixing

In this section we will show our main result, namely that the original Feynman integral (10) can be expressed as an expectation value of an observable in our topological spin foam model.

We have seen that the pure model (12) has to be gauge fixed; the complete gauge fixing is performed by choosing a admissible assignment T , and by inserting gauge fixing and Faddeev-Popov terms $K^{T_s, T_l}(l_e, s_e) = \delta_{GF}^{T_s} \delta_{GF}^{T_l} D_{FP}^{T_s} D_{FP}^{T_l}$, with $T_s = T_l = T$. We will call *observable* of the model a function $f(l_e, s_e)$ of the labels, and define its evaluation as

$$\langle f \rangle_\Delta = \frac{1}{(2\pi)^{|e|}} \int_{GF} \prod_{e \in \Delta} dl_e l_e^2 \sum_{\{s_e, \epsilon_\tau\}} f(l_e, s_e) \left(\prod_\tau \frac{e^{i\epsilon_\tau S_\tau(s_e, l_e)}}{\mathcal{V}_\tau} \right) \quad (56)$$

f is not in general gauge invariant and therefore its insertion modifies the discussion of the gauge fixing. We will be interested in particular in the class of observables that are functions of the lengths only, of a subgraph Γ drawn in Δ . The typical example is the so-called matter observable $\mathcal{O}_\Gamma(l_e) = \prod_{e \in \Gamma} G^F(l_e)$, where Γ is a subgraph of the 1-skeleton of Δ .

The introduction of such observables breaks a part of the gauge symmetries - namely the symmetry of the field l_e which acts on the vertices of Γ , turning the former gauge degrees of freedom into dynamical degrees of freedom. In the presence of matter observables the gauge fixing can no longer be symmetrical, and one needs to choose different assignments for the l 's and s 's gauge fixing. Let T_s be a maximal admissible assignment $\{\tau_v\}$, and $T_l \subset T_s$ a subset that is a maximal assignment in $\Delta \setminus \Gamma$. In other words, $T_l = \{\tau_v\}_{v \notin \Gamma}$. We then take care of the remaining

gauge symmetries by inserting the gauge fixing and Faddeev-Popov terms corresponding to these assignments. The evaluation of the observable $\mathcal{O}_\Gamma$ in the (partially) gauge fixed model reads then:

$$\langle \mathcal{O}_\Gamma \rangle_\Delta = \frac{1}{(2\pi)^{|e|}} \int \prod_{e \in \Delta} dl_e l_e^2 \sum_{\{s_e, \epsilon_\tau\}} \mathcal{O}_\Gamma(l_e) \left(\prod_\tau \frac{e^{i\epsilon_\tau S_\tau(s_e, l_e)}}{\mathcal{V}_\tau} \right) K^{T_s, T_l}(l_e, s_e) \quad (57)$$

We now suppose that Γ is a Feynman graph and Δ a triangulation of the 3-sphere S^3 such that $\Gamma \subset \Delta$. We want to show that the expectation value (57) is equal to the Feynman amplitude (10) of the graph Γ .

Recall that the triangulation Δ_k involved in (10) is a triangulation of a 3-ball, without any internal vertex nor internal edge; it is not a triangulation of S^3 . It is easy however to construct a triangulation of S^3 from Δ_k , simply by taking its *double* denoted $D\Delta_k \equiv \Delta_k \#_{S^2} \bar{\Delta}_k$, built with a copy of Δ_k in the interior of a 3-ball $B_3 \subset S^3$ glued to a copy of Δ_k in the exterior $S^3 \setminus B_3$.

The proof proceeds in three steps. Primarily, since both $D\Delta_k$ and Δ are triangulations of S^3 which contain vertices of the graph Γ , the two triangulations can be constructed out of each other by a sequence of Pachner moves that do not remove the vertices of Γ . Secondly, $\langle \mathcal{O}_\Gamma \rangle_\Delta$ is invariant under these moves, provided that the variable l_e of each edge $e \in \Gamma$ that is removed by a (2, 3) move is replaced by its value ‘on shell’, namely the one fixed by the constraint $\omega_e^\epsilon = 0$. The justification comes directly from the analysis of section III C. One can therefore write

$$\langle \mathcal{O}_\Gamma \rangle_\Delta = \langle \tilde{\mathcal{O}}_\Gamma \rangle_{D\Delta_k}, \quad \text{with} \quad \tilde{\mathcal{O}}_\Gamma = \mathcal{O}_\Gamma(l_e, l_e^\epsilon(l_e)). \quad (58)$$

The quantities $l_e^\epsilon(l_e)$ are by construction the Euclidean distances, in any embedding¹⁰ of $D\Delta_k$ in $\mathbb{R}^3$, between the vertices of Γ that are not connected by the edges of the triangulation.

Thirdly, $\tilde{I}_\Gamma \equiv \langle \tilde{\mathcal{O}}_\Gamma \rangle_{D\Delta_k}$ turns out to coincide with the Feynman amplitude I_Γ given by (10). In order to see this, let us express the gauge fixing terms for the triangulation $D\Delta_k$. We need to choose a maximal assignment T_s of $D\Delta_k$ and a assignment $T_l \subset T_s$ maximal in $D\Delta_k \setminus \Gamma$. Notice first that, since all the vertices of $D\Delta_k$ belong to the graph Γ , T_l is empty. Next, recall that, as emphasized in part II, the triangulation Δ_k of the ball is dual to a tree T ; we can take this tree to be a gauge fixing tree T_s , according to the procedure described in III D. Vertices, edges and simplices of T_s are the vertices, edges and simplices of $\Delta_k \subset D\Delta_k$. Therefore the gauge fixing is performed by inserting

$$\delta_{GF}^{T_s} = \prod_{e \in \Delta_k} (2\pi) \delta_{s_e, s_e^\epsilon} \prod_{v \in \Delta_k} \chi(\Omega_v^\epsilon) \quad ; \quad D_{FP}^{T_s} = \frac{\prod_{\tau \in \Delta_k} \mathcal{V}_\tau}{\prod_{e \in \Delta_k} l_e} \quad (59)$$

Since Δ_k has no internal vertex or edge, Δ_k and $D\Delta_k$ possess the same number of vertices and edges, all located on the boundary of the ball B , whereas the number of tetrahedra in $D\Delta_k$ is $2k$. For each tetrahedron τ in the interior of B , there exists a copy of τ in the exterior of B , sharing its edges with τ , and consequently having the same volume. Moreover, the tetrahedron τ and its copy have independent orientations ϵ_τ and ϵ'_τ . Therefore, using (59), $\tilde{I}_\Gamma$ can be written as a quantity computed on the triangulation Δ_k

$$\tilde{I}_\Gamma = \int \prod_{e \in \Delta_k} l_e dl_e \sum_{\epsilon, \epsilon' \in \{\pm 1\}^k} \mathcal{O}_\Gamma(l_e, l_{e'}^{\epsilon, \epsilon'}(l_e)) \left(\prod_{\tau \in \Delta_k} \frac{1}{\mathcal{V}_\tau} \right) e^{i \sum_e s_e^\epsilon \omega_e^{\epsilon, \epsilon'}} \prod_{v \in \Delta_k} \chi(\Omega_v^{\epsilon, \epsilon'}) \quad (60)$$

¹⁰ Note that $D\Delta_k$ can always be embedded in $\mathbb{R}^3$.

where deficit angles are given by

$$\omega_e^{\epsilon, \epsilon'}(l_e) = \sum_{\substack{\tau \in \Delta_k \\ \tau \supset e}} (\epsilon_\tau + \epsilon'_\tau) \theta_e^\tau \quad (61)$$

Finally it's easy to convince oneself that the total algebraic solid angles seen from vertices v vanish modulo 4π only if $\epsilon_\tau + \epsilon'_\tau = 0$ for every tetrahedron of Δ_k . Therefore $\tilde{I}_\Gamma = I_\Gamma$.

IV. ALGEBRAIC STRUCTURE

In order to complete our investigation, we now want to give an algebraic interpretation of the topological state sum model introduced before. This will show that this model is really a spin foam model, namely constructed purely in terms of algebraic data.

A. A Poincaré model

1. The result

We first introduce some notations. We define

$$\rho \equiv (l_e, s_e), \quad \int d\mu(\rho_e) \equiv \frac{1}{2\pi} \sum_{s_e} \int dl_e l_e^2 \quad (62)$$

and introduce a symbol associated to the tetrahedron τ with vertices 0, 1, 2, 3:

$$\left\{ \begin{array}{ccc} \rho_{23} & \rho_{13} & \rho_{12} \\ \rho_{01} & \rho_{02} & \rho_{03} \end{array} \right\} \equiv \sum_{\epsilon} \frac{e^{i\epsilon_\tau S_\tau(s_{IJ}, l_{IJ})}}{\mathcal{V}(l_{IJ})} = 2 \frac{\cos S_\tau(s_{IJ}, l_{IJ})}{\mathcal{V}(l_{IJ})}, \quad (63)$$

with

$$S_\tau(l_e, s_e) = \sum_{e \in \tau} s_e \theta_e^\tau(l_e). \quad (64)$$

The previous model can be written as a Ponzano-Regge like model [19]

$$Z_\Delta = \int \prod_e d\mu(\rho_e) \prod_\tau \left\{ \begin{array}{ccc} \rho_{e_1^\tau} & \rho_{e_2^\tau} & \rho_{e_3^\tau} \\ \rho_{e_4^\tau} & \rho_{e_5^\tau} & \rho_{e_6^\tau} \end{array} \right\}, \quad (65)$$

where the integral is over all edge labels and where e_i^τ label the 6 edges of τ .

The topological invariance of this model is essentially due to the identity (30), which can be rewritten in terms of this symbol as a pentagonal or Biedenharn-Elliott identity

$$\left\{ \begin{array}{ccc} \rho_1 & \rho_2 & \rho_3 \\ \rho_4 & \rho_5 & \rho_6 \end{array} \right\} \left\{ \begin{array}{ccc} \rho_1 & \rho_2 & \rho_3 \\ \rho'_4 & \rho'_5 & \rho'_6 \end{array} \right\} = \int d\mu(\rho) \left\{ \begin{array}{ccc} \rho_5 & \rho_1 & \rho_6 \\ \rho'_6 & \rho & \rho'_5 \end{array} \right\} \left\{ \begin{array}{ccc} \rho_6 & \rho_2 & \rho_4 \\ \rho'_4 & \rho & \rho'_6 \end{array} \right\} \left\{ \begin{array}{ccc} \rho_5 & \rho_3 & \rho_4 \\ \rho'_4 & \rho & \rho'_5 \end{array} \right\}. \quad (66)$$

All these considerations clearly suggest an interpretation of this symbol as a $6j$ -symbol for a classical group. This is indeed the case, according to the following

Theorem: $\rho = (l, s)$ labels the unitary representation of the 3-dimensional (Euclidean) Poincaré group $\text{ISO}(3)$ with mass l , and spin s ; $d\mu(\rho)$ is the Plancherel measure which arises in

the decomposition of $L^2(\text{ISO}(3))$, and the symbol $\{\rho_i\}$ defined in (63) is the normalized $6j$ -symbol of $\text{ISO}(3)$.

As a first corollary of this theorem, we get an other independent and non geometric proof of the pentagon identity (30), since this is a direct consequence of recoupling theory. An other corollary of this theorem is the fact that this $6j$ -symbol satisfies the orthogonality relation

$$\int d\mu(\rho_{12}) \left\{ \begin{matrix} \rho_{23} & \rho_{13} & \rho_{12} \\ \rho_{01} & \rho_{02} & \rho_{03} \end{matrix} \right\} \left\{ \begin{matrix} \rho_{23} & \rho_{13} & \rho_{12} \\ \rho_{01} & \rho_{02} & \rho'_{03} \end{matrix} \right\} = 2\pi \delta_{s_{03}, s'_{03}} \frac{\delta(l_{03} - l'_{03})}{l_{03}^2}. \quad (67)$$

It is interesting to give a direct and independent geometric proof of this identity. We consider two tetrahedra τ, τ' with vertices $0, 1, 2, 3$, with edge lengths and spin labels $(l_{IJ}, s_{IJ}), (l'_{IJ}, s'_{IJ})$, these labels being such that $l_{IJ} = l'_{IJ}, s_{IJ} = s'_{IJ}$ except for $(IJ) = (03)$. We want to compute

$$I = \frac{1}{2\pi} \sum_{s_{12}} \int dl_{12} l_{12}^2 \frac{\sum_{\epsilon} e^{i\epsilon S_{\tau}}}{\mathcal{V}_{\tau}} \frac{\sum_{\epsilon'} e^{i\epsilon' S_{\tau'}}}{\mathcal{V}_{\tau'}} \quad (68)$$

$$= \int dl_{12} l_{12}^2 \sum_{\epsilon, \epsilon'} \frac{\delta(\epsilon \theta_{12} - \epsilon' \theta'_{12})}{\mathcal{V}_{\tau} \mathcal{V}_{\tau'}} \left(\prod_{(ab) \neq (12)} e^{i(\epsilon s_{ab} \theta_{ab} - \epsilon' s'_{ab} \theta'_{ab})} \right) \quad (69)$$

where θ_{ij} denotes the dihedral angle at the edge (ij) . One can now use (A14)

$$\frac{\partial \theta_{03}}{\partial l_{12}} = \frac{l_{12} l_{03}}{\mathcal{V}_{\tau}}, \text{ and } \frac{\partial \theta_{12}}{\partial l_{03}} = \frac{l_{12} l_{03}}{\mathcal{V}_{\tau}} \quad (70)$$

to express the change of integration variables

$$dl_{12} l_{12} = d\theta_{03} \frac{\mathcal{V}_{\tau}}{l_{03}} \quad (71)$$

and

$$\delta(\epsilon \theta_{12} - \epsilon' \theta'_{12}) = \delta_{\epsilon, \epsilon'} \delta(l_{03} - l'_{03}) \left| \frac{\partial \theta_{12}}{\partial l_{03}} \right| = \delta_{\epsilon, \epsilon'} \delta(l_{03} - l'_{03}) \frac{\mathcal{V}_{\tau}}{l_{12} l_{03}} \quad (72)$$

Since $l_{03} = l'_{03}$, all the volume factors in the integral cancel out; and since $s_{ab} = s'_{ab}$ for $(ab) \neq (03)$, the integral reduces to

$$I = \frac{\delta(l_{03} - l'_{03})}{l_{03}^2} \int_0^{\pi} d\theta_{03} \sum_{\epsilon} e^{i\epsilon \theta_{03} (s_{03} - s'_{03})} = \frac{\delta(l_{03} - l'_{03})}{l_{03}^2} 2\pi \delta_{s_{03}, s'_{03}} \quad (73)$$

which is the desired result.

2. Proof of the Theorem

We will need some definitions. We start by describing the representation spaces of the Poincaré group. Let us fix a frame $(\vec{e}_x, \vec{e}_y, \vec{e}_z)$ of $\mathbb{R}^3$. The carrier space corresponding to a representation¹¹ $\rho \equiv (l, s)$ of $\text{ISO}(3) = \text{SO}(3) \ltimes \mathbb{R}^3$ is given by:

$$\mathcal{E}_{\rho} = \{\phi \in \mathcal{L}^2(\text{SU}(2)) / \forall \xi \in [-\pi, \pi], \phi(xh_{\xi}) = e^{is\xi} \phi(g)\}, \quad (74)$$

¹¹ The spin $s \in \frac{1}{2}\mathbb{Z}$ label representations of the little group $U(1)$. In the following we restrict to the case where all spins are integers.

where $h_\xi = e^{i\xi\sigma_3/2}$ are elements of the $U(1)$ -subgroup of rotations with axis z and angle ξ . The action of a rotation $g \in SU(2)$ and a translation $\vec{a} \in \mathbb{R}^4$ on this space is then

$$(g, 0) \phi(x) = \phi(g^{-1}x), \quad (1, \vec{a}) \phi(x) = e^{i\vec{P}_x \cdot \vec{a}} \phi(x) \quad (75)$$

where $\vec{P}_x \equiv x \cdot \vec{l}$ is the image, by the rotation associated to x , of a fixed vector $\vec{l} = le_z$ of length l .

We now define the Clebsh-Gordan coefficients. The tensor product $(l_1, s_1) \otimes (l_2, s_2)$ decomposes on representations $\rho_3 = (l_3, s_3)$ such that l_1, l_2, l_3 satisfy triangular inequalities and such that the sum of s_i 's is an integer

$$|l_1 - l_2| \leq l_3 \leq l_1 + l_2, \quad s_1 + s_2 + s_3 \in \mathbb{Z}$$

Given three lengths satisfying these inequalities, one can construct an oriented triangle in the (x, z) plane such that

$$\vec{l}_3 = a_{\theta_{13}} \cdot \vec{l}_1 + a_{\theta_{23}}^{-1} \cdot \vec{l}_2 \quad (76)$$

where one introduces the 2×2 matrix representing a rotation of angle θ and axis y

$$a_\theta = \begin{pmatrix} \cos \theta/2 & \sin \theta/2 \\ -\sin \theta/2 & \cos \theta/2 \end{pmatrix}.$$

The angles $\theta_{13}, \theta_{23} \in [0, \pi]$ are the angles between the edges (1) and (3) (resp. (2), (3)) within the oriented triangle [123] whose edges lengths are l_1, l_2, l_3 . They are characterized by

$$l_1^2 = l_2^2 + l_3^2 - 2l_2l_3 \cos \theta_{23} \quad (77)$$

and similarly for θ_{13} . The Clebsh-Gordan coefficient:

also called $3j$ -symbol, is defined (up to a normalization factor) to be the invariant map $\mathcal{E}_{l_i, s_i} \otimes \mathcal{E}_{l_j, s_j} \rightarrow \mathcal{E}_{l_k, s_k}$. The structure of this object is similar to the Clebsh-Gordan coefficient of the quantum double $\mathcal{DSU}(2)$, whose construction is described in [20]. When properly normalized, it can be written as

$$C_{x_1 x_2 x_3}^{\rho_1 \rho_2 \rho_3} = \left(\frac{l_1 l_2 l_3}{\pi} \right)^{-\frac{1}{2}} \int_{U(1)} \prod_{i=1}^3 (d\zeta_i e^{is_i \zeta_i}) \delta_{x_1 h_{\zeta_1}^{-1}} (x_3 h_{\zeta_3} a_{\theta_{13}}) \delta_{x_2 h_{\zeta_2}^{-1}} (x_3 h_{\zeta_3} a_{\theta_{23}}^{-1}) \quad (78)$$

where $d\zeta$ is the normalized Haar measure on $U(1)$. It is clear that this coefficient is such that $C_{x_1 h_{\zeta_1}^{-1} x_2 h_{\zeta_2}^{-1} x_3 h_{\zeta_3}}^{\rho_1 \rho_2 \rho_3} = C_{x_1 x_2 x_3}^{\rho_1 \rho_2 \rho_3} e^{-i(s_1 \zeta_1 + s_2 \zeta_2 - s_3 \zeta_3)}$. Moreover, it is straightforward to check (thanks to (76)) that it defines an invariant tensor, that is, $C_{g x_i}^{\rho_i} = C_{x_i}^{\rho_i}$, and that $\sum_i x_i \cdot \vec{l}_i = 0$. The condition that imposes the sum of the spins to be an integer comes from the identities $C_{x_i}^{\rho_i} = C_{-x_i}^{\rho_i} = C_{x_i h_{2\pi}}^{\rho_i} = e^{2i\pi \sum_i s_i} C_{x_i}^{\rho_i}$.

The normalization factor is chosen in order to have the resolution of the identity [3]:

$$\int d\mu(\rho_3) \left(\int dx_3 C_{x_1 x_2 x_3}^{\rho_1 \rho_2 \rho_3} \overline{C_{z_1 z_2 x_3}^{\rho_1 \rho_2 \rho_3}} \right) = \bar{\delta}_{z_1}^{s_1}(x_1) \bar{\delta}_{z_2}^{s_2}(x_2) \quad (79)$$

where the ‘bar’ denotes the complex conjugate, $d\mu(\rho_3)$ denotes the Plancherel measure (62), and

$$\bar{\delta}_{z_i}^{s_i}(x_i) = \int_{U(1)} d\zeta e^{-is_i \zeta} \delta_{z_i h_\zeta}(x_i)$$

It is worth mentioning two further properties that the Clebsh-Gordan (78) satisfies. The first one is a ‘braiding’ property which describe the change of the Clebsh-Gordan under the interchange of the labels i and j :

$$C_{x_2 x_1 x_3}^{\rho_2 \rho_1 \rho_3} = (-1)^{s_1 + s_2 - s_3} C_{x_1 x_2 x_3}^{\rho_1 \rho_2 \rho_3}, \quad (80)$$

derived thanks to the identity $h_\pi a_\theta h_{-\pi} = a_{-\theta}$. The second property expresses the dual intertwiner $\tilde{C}^{\rho_1 \rho_2 \rho_3} : \mathcal{E}_{\rho_3} \rightarrow \mathcal{E}_{\rho_1} \otimes \mathcal{E}_{\rho_2}$ in terms of the $3j$ -symbol (78) :

$$\tilde{C}^{\rho_i \rho_j \rho_k} = \overline{C^{\rho_i \rho_j \rho_k}}. \quad (81)$$

Finally, it will be important to note that, if one restricts, as we do, the spin to be integers, one can restrict the integral in (78) to be over $\xi \in [0, 2\pi]$, provided that one takes the $SO(3)$ delta function in (78), namely $\delta_g(x) \equiv \frac{1}{2}(\delta_g(x) + \delta_{-g}(x))$.

One can now construct the $6j$ -symbol, defined as a contraction of Clebsh-Gordan coefficients

$$\int dx_0 dx_1 dx_2 dx_3 \overline{C_{x_{12} x_{02} x_{01}}^{\rho_{12} \rho_{02} \rho_{01}}} C_{x_{03} x_{13} x_{01}}^{\rho_{03} \rho_{13} \rho_{01}} C_{x_{02} x_{23} x_{03}}^{\rho_{02} \rho_{23} \rho_{03}} \equiv \left\{ \begin{matrix} \rho_{23} & \rho_{13} & \rho_{12} \\ \rho_{01} & \rho_{02} & \rho_{03} \end{matrix} \right\} C_{x_{23} x_{13} x_{12}}^{\rho_{23} \rho_{13} \rho_{12}}. \quad (82)$$

The way the contractions occur is dictated by the geometry of a tetrahedron $(0, 1, 2, 3)$ with oriented edges (IJ) . There is one Clebsh-Gordan attached to each face of the tetrahedron, and one representation for each edge.

Let us now prove the theorem, by first noting that each Clebsh-Gordan of the LHS of (82) supplies two δ -functions on $SU(2)$, so the integral contains 6 delta functions. The integration is over three $SU(2)$ elements x_{0i} , and 6 $U(1)$ elements, denoted by ξ_{0i}, ξ'_{0i} . By performing a simple change of variables $x_{0i} \rightarrow x_{0i} h_{\xi_{0i}}$, one sees that the integrand depends only on the x_{0i} and the differences $\xi_{0i} - \xi'_{0i}$. Then the integration over the $SU(2)$ variables allows us to solve straightforwardly three of the delta functions. In order to write the result of integration in a convenient form, we assign an angle ξ_{IJ} to each edge (IJ) , $0 \leq I, J \leq 3$ of the tetrahedra [0123], and denote the corresponding group element by $h_{IJ} = h_{JI} \equiv h_{\xi_{IJ}}$. We also denote by $a_I^{JK} = a_I^{KJ}$ the rotation $a_{\theta_{(IJ)(IK)}}$ associated to the geometric angle between the edges (IJ) and (IK) in the triangle $[IJK]$. The later angles are fully determined by the lengths l_{IJ} . After integration, the LHS of (82) takes the form

$$\int \prod_{I < J} \left(d\xi_{IJ} e^{-is_{IJ} \xi_{IJ}} \right) \delta(h_{01} a_0^{12} h_{02} a_0^{23} h_{03} a_0^{31}) \delta_{x_{13} h_{13}^{-1}}(x_{12} h_{12} g_1) \delta_{x_{23} h_{23}^{-1}}(x_{12} h_{12} g_2) \quad (83)$$

times the prefactor $(\pi^{-1} l_{01} l_{02} l_{03})^{-1} (\pi^{-1} l_{12} l_{23} l_{31})^{-1/2}$, where we have defined

$$g_1 \equiv (a_1^{02})^{-1} h_{10} a_0^{13} h_{03} (a_3^{02})^{-1}, \quad g_2 \equiv (a_1^{20})^{-1} h_{10} (a_1^{03})^{-1}. \quad (84)$$

One can simplify the expression for g_1 if one remembers that the sum of the three interior angles of a flat triangle is equal to π . Using this fact for the triangles [012] and [023], we get $(a_1^{02})^{-1} = a_{-\pi} a_2^{01} a_0^{21}$ and $(a_3^{02})^{-1} = a_{-\pi} a_0^{32} a_2^{03}$. Plugging this into the expression for g_1 yields

$$g_1 = a_{-\pi} a_2^{10} \{a_0^{12} h_{01} a_0^{13} h_{03} a_0^{32} h_{02}\} h_{20}^{-1} a_2^{03} a_{-\pi} = a_{-\pi} a_2^{10} h_{20}^{-1} a_2^{03} a_{-\pi} = -a_2^{10} h_{20} a_2^{03}, \quad (85)$$

where for the last two equalities, we first have used that the term inside the bracket is 1 - this is due to the first delta function in (83)-, and then the property $a_\pi h a_\pi = -h^{-1}$.

In order to complete the proof of the theorem, we first have to find the solutions of the constraint expressed by the first δ -function in (83). These solutions of the constraint arise from the flatness of the tetrahedron, which leads to a remarkable identity at each vertex (see (B1)):

$$h_{IJ}^\epsilon a_I^{LI} h_{IL}^\epsilon a_i^{LK} h_{IK}^\epsilon a_I^{KJ} = -\epsilon \mathbf{1} \quad \forall I \quad (86)$$

$(IJKL)$ is an even permutation of (0123), $h_{IJ}^+ \equiv h_{\phi_{IJ}^e}$ is the rotation associated to the exterior dihedral angle of the edge (IJ) . Furthermore, if one considers (86) as an equation in ξ_{IJ} , then $\xi_{IJ} = \phi_{IJ}^e$, and this is the unique solution belonging to the interval $[0, 2\pi]$. Multiplying (86) by a_π on the left, and $a_{-\pi}$ on the right, has the effect to change all h into h^{-1} , while leaving the a 's unmodified; is one now multiplies the LHS by $h_{2\pi}^3 = -1$, one sees that $\xi_{IJ} = 2\pi - \phi_{IJ}^e$ is a solution of (86) with $+1$ in the RHS. These identities are carefully studied in appendix B. The bottom line is that the equation $h_{01} a_0^{12} h_{02} a_0^{23} h_{03} a_0^{31} = \pm 1$ (recall that we work with integer spin) admits two solutions when $\xi_{IJ} \in [0, 2\pi]$, namely

$$\xi_{IJ}^+ = \phi_{IJ}^e, \text{ or } \xi_{IJ}^- = 2\pi - \phi_{IJ}^e, \quad (87)$$

We denote the respective $h_{\xi_{IJ}^\pm}$ by $h_{IJ}^\pm$.

For each of these solutions, we use (86) to express the Euler decomposition of g_1 and g_2

$$g_1 = (a_1^{20})^{-1} h_{10}^\epsilon (a_1^{03})^{-1} = -\epsilon (h_{12}^\epsilon)^{-1} a_1^{23} (h_{13}^\epsilon)^{-1} \quad (88)$$

$$g_2 = -a_2^{10} h_{20}^\epsilon a_2^{03} = \epsilon (h_{21}^\epsilon)^{-1} (a_2^{31})^{-1} (h_{23}^\epsilon)^{-1} \quad (89)$$

Using this decomposition one can rescale $\xi_{ij} \rightarrow \xi_{ij} - \xi_{ij}^\epsilon$, $1 \leq i, j \leq 3$ in (83), and factorize from it the Clebsch-gordan coefficient $C_{x_{23}x_{13}x_{12}}^{\rho_{23}\rho_{13}\rho_{12}}$. The proportionality coefficient for each set of solutions is

$$\prod_{i < j} e^{-\epsilon i s_{ij} \phi_{ij}^\epsilon} \left(\frac{\pi}{l_{01} l_{02} l_{03}} \right) \int_\epsilon \left(\prod_i d\xi_{0i} e^{-i \xi_{0i}} \right) \delta(h_{01} a_0^{12} h_{02} a_0^{23} h_{03} a_0^{31}) \quad (90)$$

where the subscript ϵ in the integral reminds that we need to integrate around the solution $\xi_{0i} = \xi_{0i}^\epsilon$.

Now if we denote $g(\xi_{0i}) \equiv h_{01} a_0^{12} h_{02} a_0^{23} h_{03}$, (B14) gives the Jacobian of transformation from the Haar measure to the ξ_{0i} measure around a solution $g(\xi_{0i}) a_0^{31} = \pm 1$:

$$dg = \pi \frac{\mathcal{V}(l_{IJ})}{l_{01} l_{02} l_{03}} \prod_i d\xi_{0i} \quad (91)$$

where the volume of the tetrahedron appears. Therefore the expression in (90) can be written as

$$\prod_{i < j} e^{-\epsilon i s_{ij} \phi_{ij}^\epsilon} \left(\frac{\pi}{l_{01} l_{02} l_{03}} \right) \int \left(\prod_i d\xi_{0i} e^{-i \xi_{0i}} \right) \delta(\xi_{0i} - \xi_{0i}^\epsilon) \frac{l_{01} l_{02} l_{03}}{\pi \mathcal{V}(l_{IJ})} = \frac{\prod_{I < J} e^{-\epsilon i \phi_{IJ}^\epsilon}}{\mathcal{V}(l_{IJ})}. \quad (92)$$

The $6j$ -symbol is eventually obtained by summing the two contributions $\epsilon = \pm$. Therefore it reads

$$\left\{ \begin{matrix} \rho_{23} & \rho_{13} & \rho_{12} \\ \rho_{01} & \rho_{02} & \rho_{03} \end{matrix} \right\} = (-1)^{\sum s_{IJ}} 2^{\cos S_\tau(s_{IJ}, l_{IJ})} \frac{\mathcal{V}(l_{IJ})}{\mathcal{V}(l_{IJ})} \quad (93)$$

The exterior dihedral angles have been replaced by the interior ones $\theta^i = \pi - \phi^e$, leading to a factor $(-1)^{\sum s_{IJ}}$ in the formula.

B. Poincaré versus Ponzano-Regge

We have shown in the previous sections that a Feynman graph amplitude is the evaluation of the observable $\mathcal{O}_\Gamma(l_e)$ for a Poincaré spin-foam model. On the other hand, we know [8] that a usual Feynman graph is also the abelian limit of the evaluation of the same observable for the Ponzano-Regge model. This means that one should be able to relate in a direct way our Poincaré model with a suitable semi-classical limit of the Ponzano-Regge model.

In order to do so, let (u_e, v_e) be an edge-labelling of a triangulation Δ , with pairs of half-integers. We consider the square of the Ponzano-Regge model:

$$Z_\Delta^{PR^2} = \sum_{\{u_e, v_e\}} \prod_e d_{u_e} d_{v_e} \prod_\tau \{u\}_\tau \{v\}_\tau \quad (94)$$

where $\{u\}_\tau$ and $\{v\}_\tau$ denotes $6j$ -symbols of $SU(2)$ and $d_j = 2j + 1$. After a simple change of variables

$$u = j + s, \quad v = j - s$$

the partition function reads

$$Z_\Delta^{PR^2} = \sum_{\substack{\{j_e, s_e\} \\ |s_e| \leq j_e}} \prod_e (d_{j_e}^2 - (2s_e)^2) \prod_\tau \{j + s\}_\tau \{j - s\}_\tau \quad (95)$$

Next, let us introduce a length variable $l = d_j l_p$, where l_p is the Planck length. We would like to evaluate the limit $l_p \rightarrow 0$ of the model, in a sector where the typical value of l is kept fixed, and the spins s are bounded by a cut-off n much smaller than the typical value of j : $n \ll l/l_p$.

Interestingly, using the techniques presented in [15], asymptotics of $6j$ -symbols in (95) can easily be evaluated in this limit. We get:

$$\left\{ \frac{l}{2l_p} \pm s - \frac{1}{2} \right\}_\tau^2 \sim \frac{4l_p^3}{\pi} \frac{1}{\mathcal{V}(l_{ij})} \cos^2 \left[\sum_{i < j} \left(\frac{l_{ij}}{2l_p} \pm s_{ij} \right) \theta_{ij}^\tau + \frac{\pi}{4} \right] \quad (96)$$

for fixed values of the length l and the spin s . This suggests the following semi-classical limit for the model:

$$\frac{1}{2^{|e|}} \left(\frac{l_p^3}{\pi} \right)^{(|e| - |\tau|)} Z_\Delta^{PR^2} \sim \frac{1}{(2\pi)^{|e|}} \int \prod_{e \in \Delta} dl_e \prod_e l_e^2 \sum_{\{s_e \leq n\}} \sum_{\{\epsilon_\tau, \epsilon'_\tau\}} \prod_\tau \frac{e^{i\mathcal{S}_\tau(l_p)}}{\mathcal{V}_\tau} \quad (97)$$

where $\mathcal{V}_\tau$ is the volume of the tetrahedron τ with edge lengths l_e , and the action is given by (we write θ_e for θ_e^τ and ϵ, ϵ' for $\epsilon_\tau, \epsilon'_\tau$):

$$\mathcal{S}_\tau(l_p) = \sum_{e \in \tau} \frac{1}{2l_p} (\epsilon + \epsilon') l_e \theta_e(l_e) + (\epsilon - \epsilon') s_e \theta_e(l_e) + (\epsilon + \epsilon') \frac{\pi}{4} \quad (98)$$

We can reorganize the sum over the orientations¹²:

$$\sum_{\{\epsilon_\tau, \epsilon'_\tau\}} e^{i\mathcal{S}_\tau} = \sum_{\eta_\tau = \pm 1} e^{i\eta_\tau \left(\sum_{e \in \tau} \frac{l_e}{l_p} \theta_e + \frac{\pi}{2} \right)} + \sum_{\eta'_\tau = \pm 1} e^{i\eta'_\tau \sum_{e \in \tau} (2s_e) \theta_e} \quad (99)$$

¹² using $2 \cos \frac{a+b}{2} \cos \frac{a-b}{2} = \cos a + \cos b$.

and identify both Regge action and Poincaré action. Denoting the RHS of (97) by $\tilde{Z}_\Delta(n, l_p)$, we are interested in the quantity

$$Z_\Delta = \lim_{n \rightarrow \infty} \lim_{l_p \rightarrow 0} \tilde{Z}_\Delta(n, l_p) \quad (100)$$

We will show that Z_Δ is the partition function of our Poincaré model (12). Let us focus on the first limit $l_p \rightarrow 0$, n being fixed. The integrand of $\tilde{Z}_\Delta(n, l_p)$ is a product of sums which one can develop. The integral splits then into $2^{|\tau|}$ terms of the form

$$\frac{l^{2|K_2|}}{(2\pi)^{|e|}} \int \prod_{e \in \Delta} dl_e l_e^2 \sum_{\{s_e \leq 2n\}} \prod_{\tau} \frac{1}{\mathcal{V}_\tau} \prod_{\tau \in K_1} \left(\sum_{\eta'_\tau} e^{i\eta'_\tau \sum_{e \in K_1} s_e \theta_e} \right) \prod_{\tau \in K_2} \left(\sum_{\eta_\tau} \eta_\tau e^{i\frac{1}{l_p} \eta_\tau \sum_{e \in K_2} l_e \theta_e} \right) \quad (101)$$

K_1 and K_2 are disjoint sets covering the all set of tetrahedra; $|K_2|$ denotes the cardinal of K_2 . Note that we have made the change of variables $s_e \rightarrow 2s_e$, and then the spin are now integers. All terms such that $K_2 \neq \emptyset$ are oscillatory integrals, and therefore under stationary phase evaluation are typically of order $l_p^{|K_2|/2}$. The term corresponding to $K_2 = \emptyset$ is independent of l_p and is therefore the only term surviving when l_p goes to zero.

$$\lim_{l_p \rightarrow 0} \tilde{Z}_\Delta(n, l_p) = \frac{1}{(2\pi)^{|e|}} \int \prod_{e \in \Delta} dl_e l_e^2 \sum_{\{s_e \leq 2n\}} \sum_{\{\eta_\tau\}} \left(\prod_{\tau} \frac{e^{i\eta_\tau \sum_e s_e \theta_e}}{\mathcal{V}_\tau} \right) \quad (102)$$

Eventually, by taking the limit $n \rightarrow \infty$, we get the desired result and recover the Poincaré model in this semiclassical limit.

V. FEYNMAN DIAGRAMS ON HOMOGENEOUS SPACES

In this section we consider a scalar quantum field theory on spherical space-time. A Feynman amplitude of a graph Γ embedded in the unit 3-sphere $\mathcal{S}^3$ takes the form

$$I_\Gamma = \int_{\mathcal{S}^3} du_1 \cdots du_N \prod_{(ij) \in \Gamma} G_m(l_{ij}) \quad (103)$$

du_i is the normalized measure on the 3-sphere. The integrand is a product of propagators, which are functions of the dimensionless spherical distances $l_{ij} \in [0, \pi]$ between the vertices, and invariant under the action of the group $SO(4)$.

The results obtained for the flat case can directly be extended to the spherical one. Again, the idea is to gauge out the integration measure. The invariant measure is expressed in terms of spherical lengths and the Haar measure dU of $SO(4)$ in the same way as in section II.

$$du_1 \cdots du_{3+k} = dU \sum_{\epsilon \in \{\pm 1\}^k} \prod_{e \in \Delta_k} dl_e \sin l_e \prod_{\tau \in \Delta_k} \frac{1}{\mathcal{V}_\tau} \quad (104)$$

Δ_k is a triangulation, with spherical tetrahedra, of a 3-ball in $\mathcal{S}^3$, which possesses $3 + k$ vertices on its boundary, and which does not contain any internal edge. $\mathcal{V}_\tau$ is the square root of the Gram determinant $\det[\cos l_{ij}]$ (see appendix) associated to the tetrahedron τ .

A key identity relates the Gram determinants of the tetrahedra of a spherical 4-simplex, which has a form analogous to (14). We refer the reader to the appendix where the formula (17) and (18)

are established for spherical simplices. This leads to the emergence of the topological spin-foam model

$$Z_\Delta = \frac{1}{(2\pi)^{|\epsilon|}} \int \prod_{e \in \Delta} dl_e \sin^2 l_e \sum_{\{s_e, \epsilon_\tau\}} \left(\prod_\tau \frac{e^{i\epsilon_\tau S_\tau(s_e, l_e)}}{\mathcal{V}_\tau} \right) \quad (105)$$

The action for each tetrahedron reads

$$S_\tau = \sum_{e \in \tau} s_e \theta_e^\tau(l_e) \quad (106)$$

where θ_e^τ is the spherical interior dihedral angle of the edge e in τ . The Feynman amplitude I_Γ is then the expectation value of the observable $\mathcal{O}_\Gamma(l_e) = \prod_{e \in \Gamma} G_m(l_e)$ for the model (105) computed on any triangulation Δ of $\mathcal{S}^3$ which contains Γ as a subgraph. Analogous results for hyperbolic space are obtained by working on the hyperboloid and by replacing all the angles by hyperbolic angles.

Again, the model (105) takes the form of a Ponzano-Regge model. The weight associated to this model satisfies a pentagonal identity, and can be interpreted as a $6j$ -symbol. It is not difficult to convince oneself that the underlying group is the quantum double $\mathcal{DSU}(2)$ [20]; the proof of this statement is identical to the flat case, so we do not repeat it here.

The classical study of the action (106) differs from the flat case, since the eom non longer admit the so called Regge solutions $s_e = \alpha l_e$, unless $\alpha = 0$. Therefore the connection with usual Regge gravity seems to be lost. However, it is worth noting that these solutions can be reintroduced by adding a volume term $2 \sum_\tau \text{Vol}_\tau(l_e)$ to the action (106), where $\text{Vol}_\tau(l_e)$ is the spherical volume of the tetrahedra (which should not be confused with $\mathcal{V}_\tau$).

One can check that the model obtained is still topological, and that Feynman amplitudes can be written as evaluations of observables for this model as well. Let us introduce length variables $l_e = \sqrt{\Lambda} L_e$, where Λ is the cosmological constant. The new action can be written in terms of the geometrical data of a tetrahedron of curvature Λ

$$S_{\tau_\Lambda} = \sum_{e \in \tau_\Lambda} s_e \theta_e^{\tau_\Lambda}(L_e) + 2\Lambda^{3/2} \text{Vol}_{\tau_\Lambda}(L_e) \quad (107)$$

where τ_Λ is the image of τ by the re-scaling $l_e \rightarrow L_e = l_e/\sqrt{\Lambda}$, and $\theta_e^{\tau_\Lambda}$ is the dihedral angle of e in τ_Λ . Now using the spherical Schläfli identity for the tetrahedron τ_Λ

$$\sum_e L_e d\theta_e + 2\Lambda d\text{Vol} = 0,$$

we see that the variation of the action (107) gives Regge solution $s_e = \sqrt{\Lambda} L_e$. If we insert this solution into the action (107), we recover the Regge action describing discrete 3d gravity with cosmological constant Λ . Thus, the presence of a cosmological constant removes the degeneracy of the Regge solutions, and leads to the natural value $\alpha = \sqrt{\Lambda}$ for the scale parameter. Eventually let us remark that, since the topological model associated to the action (107) reduces to the Poincaré model (14) when $\Lambda \rightarrow 0$, we expect the associated symbol to be, again, the $6j$ -symbol of a deformation of the Poincaré group.

VI. GRAVITY AND BF THEORY

Before concluding, we would like to give some continuum explanation for the results we have obtained here. The fact that our state-sum model is built from $6j$ -symbols of the Poincaré group

shows, from standard arguments [11], that the topological theory which we have described in terms of a spin foam model is the quantization of a Poincaré BF theory. This theory can be written in terms of a one-form E and a connection A , both valued in the Poincaré algebra, whose generators J_i, P_i , $i = 0, 1, 2$ satisfy

$$[J^i, J^j] = 2\epsilon^{ijk} J_k, \quad [J^i, P^j] = 2\epsilon^{ijk} P_k, \quad [P^i, P^j] = 0, \quad (108)$$

and

$$E \equiv e^i P_i + b^i J_i, \quad A \equiv c^i P_i + \omega^i J_i. \quad (109)$$

The action is given by

$$S = \int_M \text{tr}(E \wedge F[A]) \quad (110)$$

where $F(A) = dA + A \wedge A$, and where the invariant trace is taken to be $\text{tr}(J_i P_j) = \delta_{ij}$, $\text{tr}(P_i P_j) = \text{tr}(J_i J_j) = 0$. In components this reads

$$S = \int_M (e_i \wedge R^i[\omega] + b_i \wedge d\omega c^i). \quad (111)$$

This model was introduced in [21] and recently studied in [22]. e is interpreted as the three dimensional frame field, ω as the spin connection; b, c are some topological matter fields. Its equations of motion when $b = c = 0$ have, as the only solution, the flat space-time $d_\omega e = 0$ and $R(\omega) = 0$. Following [2], one also expects the Feynman diagram observables to be related to Wilson lines observables for this theory

$$\mathcal{O}_\Gamma = \text{Pexp} \int_\Gamma \text{tr}(e_t J_0). \quad (112)$$

This result is reminiscent to the Polyakov reformulation of Feynman amplitudes as worldline integrals [23]. Such an interpretation provided a natural way, by replacing worldlines by worldsheets integrals, to deform field theory in terms of a dimensionfull parameter - leading to string theory. Analogously, our formulation of Feynman amplitudes in terms of spin foam models may provide a new way to think about consistent dimensionfull deformation of field theory structure, in order to take into account the quantum nature of background geometry.

VII. CONCLUSION

We have given in this paper a new perspective on closed Feynman diagrams in three dimensions. The main point was to show that the language of spin foam models - which was developed in order to address the problem of a background independent approach to quantum gravity - naturally arises in the context of quantum field theory, and that a careful study of Feynman amplitudes can even lead to a purely algebraic understanding of the quantum weights. It was also shown that usual field theory can be given a background independent perspective in the sense that the flat space geometry is purely encoded in terms of a choice of weights controlling the dynamics of the geometry.

This result is by itself very encouraging since it clearly shows that the language of spin foam is not unrelated to the usual language of field theory and also suggests a new way to investigate the semi-classical limit of quantum gravity. Namely, the Feynman diagram spin foam model provides a ‘zero order’ approximation of the full quantum gravity model, giving strong constraints on the

possible quantum gravity extensions. Such consistent extensions are known in 3d; but it should be clear from the analysis presented here and as shown in [1] that the results of this paper can be extended to higher dimensions.

We have addressed in this paper the case of scalar field closed Feynman diagrams. This clearly needs to be investigated further; first to the case of open diagrams but also to the case of higher spin fields. We expect in the later extension, the variable s which appears in our model, to acquire a more direct physical meaning. It would also be useful to have a deeper understanding of how usual Feynman diagrams can be understood as the expectation value of topological observables from the point of view of the continuum, as sketched in the last section.

Acknowledgments: We are grateful to the organizers of the conference Loops '05, Potsdam October 2005, where A. B. presented part of these results. We would like to thank R. Williams, T. Konopka, E. Livine and L. Smolin for useful discussions. A. B. would like to thank the Perimeter Institute for hospitality. This work is partially supported by a MENRT graduate student grant, by Eurodoc program from Région Rhône-Alpes and by the Government of Canada Award Program.

APPENDIX A: GEOMETRY OF SIMPLICIAL COMPLEXES

In this appendix we first recall some definitions and notations, and give a proof of the results (17), (18).

1. Basic definitions and discussion about orientation

We start by recalling basic features about D -dimensional simplices. An abstract (or combinatorial) D -simplex is a $D + 1$ -elements set $\{0, \dots, D\}$. Its $k + 1$ -elements subsets are called k -faces of the simplex. An embedded spherical D -simplex is defined by $D + 1$ normalized vectors $(e_0, \dots, e_D)$ in $\mathbb{R}^{D+1}$, $e_i^2 = 1$. The spherical edge lengths l_{ij} are given by

$$\cos l_{ij} = e_i \cdot e_j.$$

We denote by $(b_0, \dots, b_D)$ the dual vectors: $b_i \cdot e_j \equiv \delta_{ij}$, and we define :

$$\mathcal{V}(e_0, \dots, e_D) \equiv |\det(e_0, \dots, e_D)| \tag{A1}$$

$$G(e_0, \dots, e_D) \equiv \det(e_i \cdot e_j) = \det(\cos l_{ij}) = \mathcal{V}^2(e_0, \dots, e_D) \tag{A2}$$

$\mathcal{V}^2 = G$ is the determinant of the Gram matrix associated to this simplex. The *interior* dihedral angles are denoted by $\theta_{ij} \in [0, \pi]$ and defined by

$$b_i \cdot b_j = -|b_i| |b_j| \cos \theta_{ij}$$

In this paper we work only with the interior angles and refer to them simply as dihedral angles. θ_{ij} is the angle between the $(D - 1)$ -faces i and j (i.e the faces obtained by dropping the points i or j)¹³. Hence, they are labelled by the $(D - 2)$ -face F_{ij} opposite to i and j . In particular for $D = 3$, dihedral angles are labelled by the edges of the tetrahedron, and for $D = 4$ by the faces of the 4-simplex.

An orientation of an abstract D -simplex is a choice of an ordering $(0, \dots, D)$ of its points, up to even permutations [24]. A coloring (l_e, ϵ) of an oriented simplex is a set of numbers $l_e \in [0, \pi]$,

¹³ The exterior dihedral angle given by $\phi_{ij} \equiv \pi - \theta_{ij}$ is the angle between the normals to the $(D - 1)$ -simplices i, j .

labelling the edges of the simplex and satisfying triangular inequalities, together with a sign $\epsilon = \pm 1$. A given coloring promotes an abstract oriented simplex to an embedded spherical simplex; namely, it defines, up to rotations in $SO(D+1)$, normalized vectors $(e_0, \dots, e_D)$ - ordered according to the orientation - in $\mathbb{R}^{D+1}$, such that $\cos l_{ij} = e_i \cdot e_j$, and such that the sign of the determinant $\det(e_0, \dots, e_D)$ is ϵ . The embedding is then called *compatible* with the coloring. In the flat case, a coloring (l_e, ϵ) , with $l_e \in \mathbb{R}_+$ and $\epsilon = \pm 1$, specifies the geometry of a flat simplex $(0, \dots, D)$ in $\mathbb{R}^D$, namely a set of D vectors $\vec{l}_i$, $i = 1 \dots D$ represented by the oriented edges $(0i)$, and such that the sign of the determinant $\det(\vec{l}_1, \dots, \vec{l}_D)$ is ϵ . For instance, 4 ordered points $(0, 1, 2, 3)$, six positive numbers l_{ij} , $i, j = 0 \dots 3$ and an element $\epsilon \in \{\pm 1\}$, determine the geometry of a flat tetrahedron in $\mathbb{R}^3$.

An orientation of a D -simplex induces an orientation for each of its $(D-1)$ -faces by dropping the missing point in an even ordering of the points of the D -simplex in which this missing point is first. Now let's consider two oriented D -simplices sharing a $(D-1)$ -face. The orientations of these D -simplices are said consistent with each other if the shared face inherits opposite orientations from them. A triangulation Δ is said orientable if there exists a choice of consistent orientations for all simplices [24]; such a choice is then, by definition, an orientation of Δ . In this work we consider only orientable triangulations. For each of the triangulations we are working with, an orientation is implicitly chosen and fixed; and we don't refer to it anymore. A coloring of a triangulation is then a set $\{l_e, \epsilon_\sigma\}$, where $l_e \in [0, \pi]$ label the edges and $\epsilon_\sigma \in \{\pm 1\}$ label the simplices. Abusing terminology, the signs ϵ_σ will be called 'orientations' of the simplices.


With these definitions it is now straightforward to check the property illustrated in Fig 1, section II. Given two D -simplices σ_1 and σ_2 glued along a $(D-1)$ -face, a coloring $c_{\sigma_1 \cup \sigma_2} = \{l_e, \epsilon_{\sigma_1}, \epsilon_{\sigma_2}\}$ defines a map $\phi_{\sigma_1 \cup \sigma_2} \rightarrow \mathbb{R}^D$ such that ϕ_{σ_i} is compatible with the coloring $c_{\sigma_i} = \{l_e, \epsilon_{\sigma_i}\}$ induced on σ_i by $c_{\sigma_1 \cup \sigma_2}$. Then, if $\epsilon_{\sigma_1} = \epsilon_{\sigma_2}$, the two points opposite to the common face in each embedded D -simplex do not belong to the same half-space bounded by the hyperplane spanned by this face.

Let us give some precisions about the notion of deficit angle: we consider a set $\Sigma_F = \{\sigma \supset F\}$ of D -simplices sharing a given $(D-2)$ -face F in a colored triangulation. θ_F^σ denotes the dihedral angle of the face F in the simplex $\sigma \in \Sigma_F$. The deficit angle of the face F is a function $\omega_F^\epsilon(l_e)$ of edge labels and orientations $\epsilon \equiv \{\epsilon_\sigma, \sigma \in \Sigma_F\}$ of the D -simplices surrounding F . It is defined by

$$\omega_F^\epsilon(l_e) = \sum_{\sigma \in \Sigma_F} \epsilon_\sigma \theta_F^\sigma \quad (\text{A3})$$

Deficit angles are usually defined modulo 2π . They represent the curvature localized on $(D-2)$ -faces, in the sense of Regge calculus. While each D -simplex can be mapped in S^D (or $\mathbb{R}^D$ in the flat case), in general there is no global map, compatible with the coloring, of the complex Σ_F . The existence of such a map requires the deficit angle (A3) to vanish modulo 2π . We represent below, for $D=2$, and in the case of three triangles surrounding a vertex, different configurations for which the deficit angle at this vertex vanishes modulo 2π . The complex formed by the triangles is embedded in the Euclidean plane (in the flat case). We denote by ϵ the triplet $(\epsilon_1, \epsilon_2, \epsilon_3)$, where ϵ_j is the orientation of the triangle opposite to the point j .

We eventually define and describe some properties of the solid angles seen at vertices of a triangulation. In a $3D$ -triangulation Δ , we consider a vertex v and a tetrahedron τ to which v belongs. A $2d$ sphere surrounding v intersects τ along a spherical triangle: the angles of this triangle are the dihedral angles θ_e^τ of the edges of τ meeting at v , and its area, denoted by A_τ , is the solid angle seen at v within τ . The spherical triangles associated to all tetrahedra sharing v triangulate a surface called the *link* L_v of the vertex v . The total algebraic solid angle at v is



defined to be

$$\tilde{\Omega}_v^\epsilon = \sum_{\tau \supset v} A_\tau^{\epsilon_\tau}$$

where $A_\tau^{\epsilon_\tau} = A_\tau$ if $\epsilon_\tau = 1$ and $4\pi - A_\tau$ if $\epsilon_\tau = -1$. Note that by construction $\tilde{\Omega}_v^\epsilon = \Omega_v^\epsilon \pmod{4\pi}$, where the RHS is defined as in section III C.

We also define the quantity $\tilde{\omega}_e^\epsilon = \sum_{\tau \supset e} \tilde{\theta}_{e,\tau}^\epsilon$, where $\tilde{\theta}_{e,\tau}^\epsilon = \theta_e^\tau$ if $\epsilon_\tau = 1$, and $2\pi - \theta_e^\tau$ if $\epsilon_\tau = -1$. Note that this quantity is related to the deficit angle as $\tilde{\omega}_e^\epsilon = \omega_e^\epsilon \pmod{2\pi}$.

Using the relation between area and angles for a spherical triangle, which reads here $A_\tau = \sum_{e \supset v} \theta_e^\tau - \pi$, it is straightforward to check the following equality:

$$\frac{1}{2\pi} \left[\tilde{\Omega}_v^\epsilon + \sum_{e \supset v} (2\pi - \tilde{\omega}_e^\epsilon) \right] = \chi(L_v) \quad (\text{A4})$$

where $\chi(L) = |\tau| - |F| + |e|$ is the Euler characteristic of the link, $|\tau|$, $|F|$ and $|e|$ being the number of tetrahedra, faces and edges of Δ touching the vertex v , or equivalently the number of triangles, edges and vertices of the triangulation of the surface L_v . If Δ triangulates a manifold, every link L_v is homeomorphic to a 2d-sphere, and therefore $\chi(L_v) = 2$. The equality (A4) relates the total solid angle at v with the deficit angles of the edges meeting at v .

Lets study the case of a vertex 0 surrounded by four tetrahedra 1, 2, 3, 4. This configuration arises, for example, after a move (1, 4). First, if the deficit angle ω_{0i}^ϵ of the four edges (0*i*) vanish modulo 2π , then the Footnote 4, together with (A4), yield:

$$\Omega_0^\epsilon = 4\pi n(\epsilon), \quad \text{with} \quad n(\epsilon) = \begin{cases} \epsilon_1 & \text{if } \epsilon_i = \epsilon_j \quad \forall (i, j) \\ 0 & \text{if not} \end{cases} \quad (\text{A5})$$

In particular, $\Omega_0^\epsilon = 0 \pmod{4\pi}$. Now lets only suppose that the deficit angle ω_{04}^ϵ , with $\epsilon = (\epsilon_1, \epsilon_2, \epsilon_3)$, of the edge (04), vanishes modulo 2π . With the arguments exposed in section III C, this means that the complex of four tetrahedra 1, 2, 3, 4 can be mapped in $\mathbb{R}^3$, and therefore, there exists a value of the orientation ϵ_4^o such that the three other deficit angles vanishes modulo 2π as well, which implies $\Omega_0^{\epsilon_i, \epsilon_4^o} = 0 \pmod{4\pi}$. Now as a consequence¹⁴, one has $\Omega_0^{\epsilon_i, -\epsilon_4^o} \neq 0 \pmod{4\pi}$. This justifies the identity (38): the insertion of the function $\chi(\Omega_0^\epsilon)$ acts as a Kronecker symbol $\delta_{\epsilon_4, \epsilon_4^o}$, where $\epsilon_4^o = \eta \epsilon_4^\pm$.

¹⁴ Except for pathological configurations whose contribution is negligible.

2. On the geometry of the simplex

We work here in the context of spherical geometry. The results for flat space are obtained by taking the limit where $l_{ij} \rightarrow 0$ while ratios l_{ij}/l_{kl} are kept fixed. In this limit, G reduces to the square of the usual Euclidean volume, and we have the following correspondence :

$$\mathcal{V}(e_0, \dots, e_D) \sim D!V(l_{ij}), \quad \sin l_{ij} \sim l_{ij}, \quad \cos(l_{ij}) - 1 \sim -\frac{1}{2}l_{ij}^2. \quad (\text{A6})$$

Abusing terminology, we will call $\mathcal{V}$ the ‘volume’ of the simplex, keeping in mind that it reduces to the volume only in the flat space limit.

Next we will need the following result. Let $(e_0, \dots, e_D)$ a D -simplex and L_k be the subspace $\text{span}\{b_0, \dots, b_{k-1}\}$ generated by the vectors dual to $(e_0, \dots, e_{k-1})$. By construction L_k is the subspace orthogonal to $\text{span}\{e_k, \dots, e_D\}$. We denote by $(h_0, \dots, h_{k-1})$ the orthogonal projection of $(e_0, \dots, e_{k-1})$ onto L_k . In other words, the h_i are defined to be the vectors of $\mathbb{R}^{D+1}$ which satisfy the properties

$$h_i \cdot b_j = \delta_{ij} \quad \forall j < k, \quad \text{and} \quad h_i \cdot e_j = 0 \quad \forall j \geq k \quad (\text{A7})$$

One then has:

Proposition 1 *The following equalities hold*

$$\mathcal{V}(e_0, \dots, e_D) = \mathcal{V}(h_0, \dots, h_{k-1}, e_k, \dots, e_D) \quad (\text{A8})$$

$$= \mathcal{V}(h_0, \dots, h_{k-1}) \mathcal{V}(e_k, \dots, e_D) = \frac{\mathcal{V}(e_k, \dots, e_D)}{\mathcal{V}(b_0, \dots, b_{k-1})} \quad (\text{A9})$$

where the function $\mathcal{V}$ is defined to be $\mathcal{V}(v_1, \dots, v_n) \equiv \det(v_1, \dots, v_n)$ within $\text{span}(v_1, \dots, v_n)$.

The first equality arises from elementary column operations on the determinant, the second one by orthogonality, and the third one from the duality of the basis $(h_i)_{0 \leq i \leq k-1}$ and $(b_i)_{0 \leq i \leq k-1}$ within L_k .

If one applies this result for $k = 1$ one sees that

$$|b_i| = |h_i|^{-1}$$

where $|h_i|$ is the height of the $D - 1$ face i , defined as

$$\mathcal{V}_i = \frac{\mathcal{V}}{|h_i|}, \quad (\text{A10})$$

$\mathcal{V}$ being the D -simplex ‘volume’ and $\mathcal{V}_i$ the volume of the face i . The proposition for $k = 2$ leads to the identity

$$\mathcal{V}\mathcal{V}_{ij} = \mathcal{V}_i\mathcal{V}_j \sin \theta_{ij} \quad (\text{A11})$$

where $\mathcal{V}_{ij}$ is the ‘volume’ of the $D - 2$ face obtained by dropping i, j and θ_{ij} the dihedral angle between the faces i and j .

Using the well known relation between the cofactor matrix and the inverse matrix one easily sees that the dual vectors can be expressed through the derivative of the volume:

$$\mathcal{V}b_\mu^i = \frac{\partial \mathcal{V}}{\partial e_i^\mu}. \quad (\text{A12})$$

This implies the derivative formula

$$\frac{\partial \mathcal{V}^2}{\partial \cos l_{ij}} = -2\mathcal{V}_i \mathcal{V}_j \cos \theta_{ij}. \quad (\text{A13})$$

This is shown if one looks at the variation

$$\delta \mathcal{V}^2 = 2\mathcal{V}^2 b_\mu^i \delta e_i^\mu = 2\mathcal{V}^2 \sum_{i,j} (b^i \cdot b^j) e_j \cdot \delta e_i = 2\mathcal{V}^2 \sum_{i < j} (b^i \cdot b^j) \delta(e_j \cdot e_i) = -2\mathcal{V}_i \mathcal{V}_j \sum_{i < j} \cos \theta_{ij} \delta \cos l_{ij}$$

where summations over repeated indices is understood when it is not explicit, and where i and μ indices are raised and lowered with flat metric. From (A11, A13), one can compute the derivative

$$\frac{\partial \theta_{ij}}{\partial l_{ij}} = \frac{\mathcal{V}_{ij} \sin l_{ij}}{\mathcal{V}}. \quad (\text{A14})$$

To show this, let's start from the square of (A11) and derive it with respect to $\cos l_{ij}$. Since neither $\mathcal{V}_i, \mathcal{V}_j$ nor $\mathcal{V}_{ij}$ depends on l_{ij} , one obtains

$$\frac{\partial \mathcal{V}^2}{\partial \cos l_{ij}} \mathcal{V}_{ij}^2 = \mathcal{V}_i^2 \mathcal{V}_j^2 \frac{\partial \sin^2 \theta_{ij}}{\partial \cos l_{ij}} = (\mathcal{V}_i \mathcal{V}_j \sin \theta_{ij})(2\mathcal{V}_i \mathcal{V}_j \cos \theta_{ij}) \frac{\partial \theta_{ij}}{\partial \cos l_{ij}}.$$

Using once again the identities (A11, A13), the RHS is equal to

$$-\mathcal{V} \mathcal{V}_{ij} \frac{\partial \mathcal{V}^2}{\partial \cos l_{ij}} \frac{\partial \theta_{ij}}{\partial \cos l_{ij}} = \frac{\mathcal{V} \mathcal{V}_{ij}}{\sin l_{ij}} \left(\frac{\partial \mathcal{V}^2}{\partial \cos l_{ij}} \right) \left(\frac{\partial \theta_{ij}}{\partial l_{ij}} \right),$$

from which (A14) follows. Finally we derive a last derivative identity by taking the derivative of (A13) with respect to $\cos l_{ij}$. With the help of (A14, A11) one obtains

$$\frac{\partial^2 \mathcal{V}^2}{(\partial \cos l_{ij})^2} = -2\mathcal{V}_{ij}^2. \quad (\text{A15})$$

Next we will show the following

Proposition 2 *We consider a spherical 4-simplex $(e_0, \dots, e_4)$. We denote by l_{ij} its lengths and $G \equiv G(e_0, \dots, e_4)$ its Gram determinant. Let $\mathcal{V}_j$ be the volume and ϵ_j the orientation of the tetrahedron τ_j obtained by dropping the vertex j . Then:*

$$\left. \frac{\partial G}{\partial l_{04}} \right|_{l_{04}^\pm} = \mp 2 \sin l_{04} \mathcal{V}_0 \mathcal{V}_4 \quad (\text{A16})$$

$$\left. \frac{\partial G}{\partial \omega_{04}^\epsilon} \right|_{l_{04}^\pm} = -2\epsilon_1 \epsilon_2 \epsilon_3 \frac{\mathcal{V}_1 \mathcal{V}_2 \mathcal{V}_3}{\sin l_{04}} \quad (\text{A17})$$

where $(l_{04}^\pm, \epsilon_i)$ are solution of $G = 0$ and $\omega_{04}^\epsilon = 0 \pmod{2\pi}$, with $l_{04}^- \leq l_{04}^+$.

The first equality simply arises from (A13) and from the fact that the 4D dihedral angle of the face $[123]$ is π if $l_{04} = l_{04}^+$ and 0 if $l_{04} = l_{04}^-$.

To show the second equality, we first use the Proposition 1 for $D = 3$ and $k = 2$ to express the volume $\mathcal{V}_i$ of the tetrahedron τ_i , $i = 1, 2, 3$, as

$$\mathcal{V}_i = \mathcal{V}(h_j, h_k) \mathcal{V}(e_0, e_4) = |h_j| |h_k| \sin \theta_{ij} \sin l_{04} \quad (\text{A18})$$

where ijk is a permutation of 123, and $|h_l|$ is the height of the point l in the face $[04l]$. Indeed, first, recall that $b_j \cdot b_k = -|b_j||b_k| \cos \theta_{jk}$, where θ_{jk} is the dihedral angle between the faces $[04j]$ and $[04k]$ in τ_i ; second, writing $h_j \cdot h_k = |h_j||h_k| \cos \theta'_{ij}$, the duality of the basis (h_j, h_k) and (b_j, b_k) within the plane $\text{span}(b_j, b_k)$ implies the relation $\theta'_{ij} = \theta_{ij}$.

Now, still with the Proposition 1, the volume of the 4-simplex factorizes as

$$\mathcal{V} = \mathcal{V}(h_1, h_2, h_3) \mathcal{V}(e_0, e_4) = |h_1||h_2||h_3| \mathcal{V}(\tilde{h}_1, \tilde{h}_2, \tilde{h}_3) \sin l_{04} \quad (\text{A19})$$

Since variations of the dihedral angles leave heights $|h_i|$ and length l_{04} invariant, one only has to compute the variation of the quantity

$$\mathcal{V}^2(\tilde{h}_1, \tilde{h}_2, \tilde{h}_3) = \det(\cos \theta_{ij})$$

A first way to perform this variation is to use (A13) with the following correspondence $l_{ij} \rightarrow \theta_{ij}$, and $\theta_{ij} \rightarrow \alpha_{ij}$, where α_{ij} is the dihedral angle between edges i and j in the spherical triangle defined by $(\tilde{h}_1, \tilde{h}_2, \tilde{h}_3)$. One can convince oneself that, when $G = 0$, this spherical triangle is degenerate and, furthermore, $\cos \alpha_{ij} = -\epsilon_i \epsilon_j$. Consequently, one gets

$$\delta \mathcal{V}^2(\tilde{h}_1, \tilde{h}_2, \tilde{h}_3) = -2\epsilon_1 \epsilon_2 \epsilon_3 \sin \theta_{12} \sin \theta_{13} \sin \theta_{23} (\epsilon_1 \theta_{23} + \epsilon_2 \theta_{13} + \epsilon_3 \theta_{12}) \quad (\text{A20})$$

where the term in parenthesis is the variation of the dihedral angle ω_{04}^ϵ .

An other way to obtain this variation is a direct computation of the determinant. Indeed,

$$\delta \mathcal{V}^2(\tilde{h}_1, \tilde{h}_2, \tilde{h}_3) = -2 \sin \theta_{12} (\cos \theta_{12} - \cos \theta_{13} \cos \theta_{23}) \delta \theta_{12} + \text{perm} \quad (\text{A21})$$

Now, since the deficit angle $\omega_{04}^\epsilon = \epsilon_1 \theta_{23} + \epsilon_2 \theta_{13} + \epsilon_3 \theta_{12}$ vanishes modulo 2π , one can write

$$\begin{aligned} \cos \theta_{12} - \cos \theta_{13} \cos \theta_{23} &= \cos \epsilon_3 \theta_{12} - \cos \theta_{13} \cos \theta_{23} \\ &= \cos(\epsilon_1 \theta_{23} + \epsilon_2 \theta_{13}) - \cos \theta_{13} \cos \theta_{23} = \epsilon_1 \epsilon_2 \sin \theta_{23} \sin \theta_{13}, \end{aligned} \quad (\text{A22})$$

and insert this identity in (A21), to get (A20).

Eventually, the variation of $G = \mathcal{V}^2$ reads

$$\delta G = -2\epsilon_1 \epsilon_2 \epsilon_3 |h_1|^2 |h_2|^2 |h_3|^2 \sin^2 l_{04} \sin \theta_{12} \sin \theta_{13} \sin \theta_{23} \delta \omega_{04}^\epsilon \quad (\text{A23})$$

which, thanks to (A18), can be written as

$$\delta G = -2\epsilon_1 \epsilon_2 \epsilon_3 \frac{\mathcal{V}_1 \mathcal{V}_2 \mathcal{V}_3}{\sin l_{04}} \delta \omega_{04}^\epsilon \quad (\text{A24})$$

This last equality finally shows (A17).

APPENDIX B: FLATNESS IDENTITIES FOR THE TETRAHEDRON

We consider a flat tetrahedron $IJKL$ with edge lengths l_{ij} . We define: $\theta_I^{JK} \in [0, \pi]$ to be the geometric angle between the edges (IJ) and (IK) within the triangle IJK , $\phi_{IJ}^\epsilon \in [0, \pi]$ to be the exterior dihedral angle of the edge (IJ) . We introduce the matrices $a_I^{JK} \equiv a_{\theta_I^{JK}}$ and $h_{IJ}^+ \equiv h_{\phi_{IJ}^\epsilon}$, $h_{IJ}^- \equiv h_{2\pi - \phi_{IJ}^\epsilon}$, with

$$h_\phi = \begin{pmatrix} e^{i\phi/2} & 0 \\ 0 & e^{-i\phi/2} \end{pmatrix}, \quad a_\theta = \begin{pmatrix} \cos \theta/2 & \sin \theta/2 \\ -\sin \theta/2 & \cos \theta/2 \end{pmatrix}$$

Note that, by construction, $h_{IJ} = h_{JI}$ and $a_I^{JK} = a_I^{KJ}$. Any $SU(2)$ matrices acts as a rotation on $\mathbb{R}^3$ by adjoint action $g \cdot (x^i \sigma_i) \equiv g(x^i \sigma_i)g^{-1}$; as such h_ξ is a rotation of axis z and angle ξ , a_θ a rotation of axis y and angle θ .

Then for all even permutation $(IJKL)$ of (0123) , the following vertex identities hold:

$$h_{IJ}^\epsilon a_I^{LI} h_{IL}^\epsilon a_i^{LK} h_{IK}^\epsilon a_I^{KJ} = -\epsilon \mathbf{1} \quad (\text{B1})$$

Moreover, if one considers (B1) as an equation in ξ_{IJ} where $\xi_{IJ} \in [0, 2\pi]$ and $h_{IJ} = h_{\xi_{IJ}}$, then $\xi_{IJ} = \phi_{IJ}^e$ (the exterior dihedral angle) is the only solution for $\epsilon = +1$, and similarly $\xi_{IJ} = 2\pi - \phi_{IJ}^e$ is the only solution in $[0, 2\pi]$ of (B1) for $\epsilon = -1$.

To prove this, let us focus on the vertex 0, and let us first give a geometrical interpretation to (B1). We start from a configuration of the edges meeting at 0 such that the edge (01) is along the axis z , the edge (02) is in the plane (x, z) and in the direction of the axis x , and the edge (03) is in the direction of the axis y . After the rotation a_0^{21} , the edge (02) is along the axis z ; after the rotation h_{02}^+ the edge (03) is in the plane (x, z) . The combination $h_{02}^+ a_0^{21}$ just amounts to a cyclic permutation of the edges $(0i)$: after doing three permutations we are back where we have started. Therefore the product (B1) is equal to ± 1 .

In order to establish our statement, let's compute the two sides of the equality:

$$h_{03} a_0^{32} h_{02} = -\epsilon (a_0^{13})^{-1} (h_{01})^{-1} (a_0^{12})^{-1} \quad (\text{B2})$$

In detail the equations are

$$\begin{aligned} -\epsilon e^{i(\xi_{01} + \xi_{02} + \xi_{03})/2} \cos \theta_0^{32}/2 &= \cos \theta_0^{13}/2 \cos \theta_0^{12}/2 - e^{i\xi_{01}} \sin \theta_0^{13}/2 \sin \theta_0^{12}/2 \quad (a) \\ \epsilon e^{i(\xi_{01} + \xi_{02} - \xi_{03})/2} \sin \theta_0^{32}/2 &= \sin \theta_0^{13}/2 \cos \theta_0^{12}/2 + e^{-i\xi_{01}} \cos \theta_0^{13}/2 \sin \theta_0^{12}/2 \quad (b), \end{aligned}$$

as well as their complex conjugate. The equations (a) and (b) provide four real equations whose only three are independent, because of the unitarity condition for the matrices in (B2). Indeed, considering $a\bar{a} + a\bar{b}$, where $\bar{a}$ denotes the complex conjugate of the identity (a), leads to the trivial equality $1 = 1$. Now by taking $a\bar{a} - a\bar{b}$, we get

$$c_{32} = c_{12}c_{13} - \cos \xi_{01} s_{13}s_{12} \quad (\text{B3})$$

and two similar equations obtained by permutation of indices. c_{ij}, s_{ij} denote $\cos \theta_0^{ij}, \sin \theta_0^{ij}$. These equations in ξ_{0i} are those satisfied by the exterior dihedral angles. Consequently

$$\cos \xi_{0i} = \cos \phi_{0i}^e. \quad (\text{B4})$$

Therefore, $\xi_{0i} = \phi_{0i}^e$ or $\xi_{0i} = 2\pi - \phi_{0i}^e$, since by hypothesis $\xi_{0i} \in [0, 2\pi]$.

In order to choose between these two possibilities we need an additional geometrical interpretation of these angles: if we surround the vertex 0 by a $2d$ -sphere, the three edges meeting at 0 will intersect this sphere along three vertex of a spherical triangle, the interior of this triangle being inside the tetrahedra. θ_0^{ij} is then the spherical length of the edge (ij) of this triangle, and ϕ_{0i}^e is the exterior dihedral angle at the vertex i . The area of this spherical triangle is given by

$$A^e = 2\pi - \phi_{01}^e + \phi_{02}^e + \phi_{03}^e, \quad 0 < A < 2\pi. \quad (\text{B5})$$

We can take the imaginary part of (a), which implies

$$\epsilon \frac{\sin \xi_{01}}{\sin A/2} = \frac{\cos \theta_0^{23}/2}{\sin \theta_0^{13}/2 \sin \theta_0^{12}/2} \equiv C_1 > 0 \quad (\text{B6})$$

where $A \equiv 2\pi - \xi_{01} + \xi_{02} + \xi_{03}$. By symmetry one can get similar identities leading to positive constants C_i , $i = 1 \dots 3$, and therefore we find

$$\frac{\sin \xi_{0i}}{\sin \xi_{0j}} = \frac{C_i}{C_j} > 0 \quad \forall (ij) \quad (\text{B7})$$

This last property imposes all the ξ_{0i} to coincide to ϕ_{0i}^e or all to $2\pi - \phi_{0i}^e$. In the first case we have $A = A^e$; in the second $A = -2\pi - A^e$; in both cases $\sin A/2 > 0$. Therefore one sees that the sign of $\sin \xi_{01}$ is ϵ . This shows that the solutions are $\phi_{0i} = \phi_{0i}^e$ when $\epsilon = +1$ and $\phi_{0i} = 2\pi - \phi_{0i}^e$ when $\epsilon = -1$.

As an application of the calculation of (B2), let us show the expression (91) for the jacobian. By considering the Euler decomposition of the matrix $a_0^{12} h_{02} a_0^{23} \equiv h_\phi a_\theta h_\psi$ we write

$$g(\xi_{0i}) = h_{01} h_\phi a_\theta h_\psi h_{03} \quad (\text{B8})$$

The Euler parameters do not depend on ξ_{01}, ξ_{02} . Therefore

$$dg = \frac{1}{2} \sin \theta d\theta d(\xi_{03} + \phi) d(\xi_{02} + \psi) \quad (\text{B9})$$

$$= \frac{1}{2} \sin \theta d\theta \frac{d\xi_{03}}{2\pi} \frac{d\xi_{02}}{2\pi}. \quad (\text{B10})$$

Now we know from previous calculations that the angles ξ_{01} and θ are related to each other in the following way

$$\cos \theta = c_{12} c_{23} - \cos \xi_{01} s_{12} s_{23} \quad (\text{B11})$$

where c_{ij}, s_{ij} denotes $\cos \theta_0^{ij}, \sin \theta_0^{ij}$. Consequently

$$\sin \theta d\theta = 2\pi \sin \xi_{01} s_{12} s_{13} \frac{d\xi_{01}}{2\pi} \quad (\text{B12})$$

$$= 2\pi \frac{\mathcal{V}(l_{ij})}{l_{01} l_{02} l_{03}} \frac{d\xi_{01}}{2\pi} \quad (\text{B13})$$

$\mathcal{V}(l_{ij})$ is (3! times) the volume of the tetrahedron whose edge lengths are l_{ij} . The last formula holds as long as ξ_{01} is a solution of (B1), since this angle is then equal, up to a sign and modulo 2π , to the exterior dihedral angle of the edge (01). Hence,

$$dg = \pi \frac{\mathcal{V}(l_{ij})}{l_{01} l_{02} l_{03}} \prod_i \frac{d\xi_{0i}}{2\pi} \quad (\text{B14})$$

-
- [1] A. Baratin, L. Freidel *Hidden Quantum Gravity in 4d Feynman diagrams: Emergence of spin foams*, To appear.
 - [2] L. Freidel and D. Louapre, "Ponzano-Regge model revisited. I: Gauge fixing, observables and interacting spinning particles," *Class. Quant. Grav.* **21**, 5685 (2004) [arXiv:hep-th/0401076].
 - [3] L. Freidel and D. Louapre, "Ponzano-Regge model revisited. II: Equivalence with Chern-Simons," arXiv:gr-qc/0410141.
 - [4] J. W. Barrett, "Geometrical measurements in three-dimensional quantum gravity," *Int. J. Mod. Phys. A* **18S2**, 97 (2003) [arXiv:gr-qc/0203018].
 - [5] J. W. Barrett, J. M. Garcia-Islas and J. F. Martins, "Observables in the Turaev-Viro and Crane-Yetter models," arXiv:math.qa/0411281.

- [6] J. W. Barrett, “Feynman loops and three-dimensional quantum gravity,” *Mod. Phys. Lett. A* **20**, 1271 (2005) [arXiv:gr-qc/0412107].
- [7] K. Noui and A. Perez, “Three dimensional loop quantum gravity: Coupling to point particles,” *Class. Quant. Grav.* **22**, 4489 (2005) [arXiv:gr-qc/0402111].
- [8] L. Freidel and E. R. Livine, “Ponzano-Regge model revisited. III: Feynman diagrams and effective field theory,” arXiv:hep-th/0502106.
- [9] L. Freidel and E. R. Livine, “Effective 3d quantum gravity and non-commutative quantum field theory,” arXiv:hep-th/0512113.
- [10] L. Freidel and S. Majid, “Noncommutative harmonic analysis, sampling theory and the Duflo map in 2+1 quantum gravity,” arXiv:hep-th/0601004.
- [11] J. C. Baez, “Spin foam models,” *Class. Quant. Grav.* **15**, 1827 (1998) [arXiv:gr-qc/9709052]. L. Freidel and K. Krasnov, “Spin foam models and the classical action principle,” *Adv. Theor. Math. Phys.* **2**, 1183 (1999) [arXiv:hep-th/9807092]. D. Oriti, “Spacetime geometry from algebra: Spin foam models for non-perturbative quantum gravity,” *Rept. Prog. Phys.* **64**, 1489 (2001) [arXiv:gr-qc/0106091]. A. Perez, “Spin foam models for quantum gravity,” *Class. Quant. Grav.* **20**, R43 (2003) [arXiv:gr-qc/0301113].
- [12] L. Freidel and A. Starodubtsev, *Quantum gravity in terms of topological observables*, arXiv:hep-th/0501191.
- [13] J. W. Barrett, “Feynman diagrams coupled to three-dimensional quantum gravity,” *Class. Quant. Grav.* **23**, 137 (2006) [arXiv:gr-qc/0502048].
- [14] I.G. Korepanov, *Invariants of PL Manifolds from Metrized simplicial Complexes. Three-Dimensional Case*, *J. Nonlin. Math. Phys* **8**, (2001) 196-210.
- [15] L. Freidel, D. Louapre *Asymptotics of 6j and 10j symbols*, *Class. Quant. Grav.* **20**, 1267-1294, (2003) [arXiv:hep-th/0209134].
- [16] Regge T, *General Relativity without Coordinates*, *Nuovo Cim. A* **19** (1961), 558-571.
- [17] Yuka U. Taylor and Christopher T. Woodward, *Spherical tetrahedra and invariants of 3-manifolds*, arXiv:math.GT/0406228.
- [18] L. Freidel, D. Louapre *Diffeomorphisms and spin foam model*, *Nucl. Phys. B* **662**, 279-298 (2003) [arXiv:gr-qc/0212001].
- [19] G. Ponzano, T. Regge *Semi-classical limit of Racah coefficients* in *Spectroscopic and Group theoretical methods in physics*, ed F Bloch, North-holland, New-York (1968).
- [20] T.H. Koornwinder, F.A. Bais and N.M. Muller, *Tensor product representations of the quantum double of a compact group*, *Commun. Math. Phys.* **198**, 157 (1998) [arXiv:q-alg/9712042]
- [21] S. Carlip, J. Gegenberg, *Gravitating topological matter in 2+1 dimensions*, *Phys. Rev. D* **44**, 424 (1991)
- [22] L. Freidel, R.B. Mann, Eugeniu M. Popescu, *Canonical analysis of the BCEA topological matter model coupled to gravitation in 2+1 dimensions*, *Class. Quant. Grav.* **22** 3363-3380 (2005) [arXiv:gr-qc/0411117].
- [23] A. M. Polyakov (Moscow, ITEP), *Gauge Fields and Strings*, Chur, Switzerland: Harwood (1987) (Contemporary Concepts in Physics, 3) 301p.
- [24] Singer, I.M, and Thorpe, J.A. *Lecture Notes on Elementary Geometry and Topology*, New-York: Springer-Verlag (1996).